

2004
Update



India

EPIDEMIOLOGICAL FACT SHEETS
ON HIV/AIDS AND SEXUALLY TRANSMITTED INFECTIONS

HIV/AIDS estimates

In 2003 and during the first quarter of 2004, UNAIDS and WHO worked closely with national governments and research institutions to recalculate current estimates on people living with HIV/AIDS. These calculations are based on the previously published estimates for 1999 and 2001 and recent trends in HIV/AIDS surveillance in various populations. A methodology developed in collaboration with an international group of experts was used to calculate the new estimates on prevalence and incidence of HIV and AIDS deaths, as well as the number of children infected through mother-to-child transmission of HIV. Different approaches were used to estimate HIV prevalence in countries with low-level, concentrated or generalised epidemics. The current estimates do not claim to be an exact count of infections. Rather, they use a methodology that has thus far proved accurate in producing estimates that give a good indication of the magnitude of the epidemic in individual countries. However, these estimates are constantly being revised as countries improve their surveillance systems and collect more information.

Adults in this report are defined as women and men aged 15 to 49. This age range covers people in their most sexually active years. While the risk of HIV infection obviously continues beyond the age of 50, the vast majority of those who engage in substantial risk behaviours are likely to be infected by this age. The 15 to 49 range was used as the denominator in calculating adult HIV prevalence.

Estimated number of adults and children living with HIV/AIDS, end of 2003

These estimates include all people with HIV infection, whether or not they have developed symptoms of AIDS, alive at the end of 2003:

Adults and children	5,100,000		
Low estimate	2,500,000		
High estimate	8,500,000		
Adults (15-49)	5,000,000	Adult rate (%)	0.9
Low estimate	2,500,000	Low estimate	0.5
High estimate	8,200,000	High estimate	1.5
Children (0-15)	120,000		
Low estimate	55,000		
High estimate	260,000		
Women (15-49)	1,900,000		
Low estimate	710,000		
High estimate	2,400,000		

Estimated number of deaths due to AIDS

Estimated number of adults and children who died of AIDS during 2003:

Deaths in 2003	0
Low estimate	0
High estimate	0

Estimated number of orphans

Estimated number of children who have lost their mother or father or both parents to AIDS and who were alive and under age 17 at the end of 2003:

Current living orphans	0
Low estimate	0
High estimate	0

Assessment of the epidemiological situation 2004

The evidence of HIV was first documented in Chennai in southern India in 1986. From then until March 2004, 68,809 AIDS cases have been reported to the National AIDS Control Organization. Heterosexual route is the predominant mode of transmission, followed by injecting drug use.

Nationwide, annual HIV sentinel surveillance (HSS) was started in 1998 and so far six rounds have been completed. The numbers of ANC sentinel sites have increased considerably over the years, however, the vulnerable population groups such as MSMs, IDUs and CSWs remain largely under-represented.

The national adult HIV prevalence is 0.8%. However, the aggregate data mask the multiple and diverse HIV sub-epidemics in the country. Of the 35 States of India, 6 States, four in southern India (Andhra Pradesh, Tamil Nadu, Maharashtra, Karnataka) and two in north eastern India (Manipur and Nagaland) have generalized epidemics with HIV prevalence rate of above 1% among pregnant women. These six States account for nearly 80% of all reported AIDS cases in the country. In the southern States, heterosexual transmission accounts for majority of the reported cases, whereas in the north-eastern States of Manipur and Nagaland, injecting drug use is the predominant mode of transmission, where the prevalence among injecting drug users in 2003 was 56%. More than 50% of the CSWs in urban southern States are infected.

Three States/Union Territories (Gujarath, Goa and Pondicherry) have concentrated epidemics and the rest of the states in India have low level epidemics. In the six States with generalized epidemics, the HIV prevalence among pregnant women has remained constant over the last three years, indicating that the epidemic may have stabilized.

Although the overall prevalence of HIV is below 1%, due to the large population size, India has a large number of people living with HIV/AIDS, second only to South Africa. By the end of 2002, there were an estimated 3.8-4.6 million PLWHAs in India. Of these, 600,000 PLWHAs are in need of antiretroviral therapy but less than 30,000 are currently receiving ART. The Government of India launched the national antiretroviral programme in April 2004. The national ART programme will be scaled up as human and financial resources become available.

With a high prevalence of TB infection in India, the problem of TB related to HIV infection also poses a major public health challenge, particularly in the high HIV burden States. The exact magnitude of HIV among TB patients is unknown as routine surveillance of HIV among TB patients is not undertaken.

UNAIDS/WHO Working Group on Global HIV/AIDS and STI Surveillance

Global Surveillance of HIV/AIDS and sexually transmitted infections (STIs) is a joint effort of WHO and UNAIDS. The UNAIDS/WHO Working Group on Global HIV/AIDS and STI Surveillance, initiated in November 1996, guides respective activities. The primary objective of the Working Group is to strengthen national, regional and global structures and networks for improved monitoring and surveillance of HIV/AIDS and STIs. For this purpose, the Working Group collaborates closely with national AIDS programmes and a number of national and international experts and institutions. The goal of this collaboration is to compile the best information available and to improve the quality of data needed for informed decision-making and planning at national, regional, and global levels. The Epidemiological Fact Sheets are one of the products of this close and fruitful collaboration across the globe.

Within this framework, the Fact Sheets collate the most recent country-specific data on HIV/AIDS prevalence and incidence, together with information on behaviours (e.g. casual sex and condom use) which can spur or stem the transmission of HIV.

Not unexpectedly, information on all of the agreed upon indicators was not available for many countries in 2003. However, these updated Fact Sheets do contain a wealth of information which allows identification of strengths in currently existing programmes and comparisons between countries and regions. The Fact Sheets may also be instrumental in identifying potential partners when planning and implementing improved surveillance systems.

The fact sheets can be only as good as information made available to the UNAIDS/WHO Working Group on Global HIV/AIDS and STI Surveillance. Therefore, the Working Group would like to encourage all programme managers as well as national and international experts to communicate additional information to them whenever such information becomes available. The Working Group also welcomes any suggestions for additional indicators or information proven to be useful in national or international decision-making and planning.

Basic indicators

For consistency reasons the data used in the table below are taken from official UN publications.

DEMOGRAPHIC DATA	YEAR	ESTIMATE	SOURCE
Total population (thousands)	2004	1,081,229	UN population division database
Female population aged 15-24 (thousands)	2004	98663	UN population division database
Population aged 15-49 (thousands)	2004	564944	UN population division database
Annual population growth rate (%)	1992-2002	1.8	UN population division database
% of urban population	2003	28.2	UN population division database
Average annual growth rate of urban population	2000-2005	2.28	UN population division database
Crude birth rate (births per 1,000 pop.)	2004	23.1	UN population division database
Crude death rate (deaths per 1,000 pop.)	8.3	8.3	UN population division database
Maternal mortality rate (per 100,000 live births)	2000	540	WHO (WHR2004)/UNICEF
Life expectancy at birth (years)	2002	61	World Health Report 2004, WHO
Total fertility rate	2002	3.1	World Health Report 2004, WHO
Infant mortality rate (per 1,000 live births)	2000	77	World Health Report 2004, WHO
Under 5 mortality rate (per 1,000 live births)	2000	96	World Health Report 2004, WHO

SOCIO-ECONOMIC DATA	YEAR	ESTIMATE	SOURCE
Gross national income, ppp, per capita (Int.\$)	2002	80	World Health Report 2004, WHO
Gross domestic product, per capita % growth	2001-2002	2.8	World Bank
Per capita total expenditure on health (Int.\$)	2001	80	World Health Report 2004, WHO
General government expenditure on health as % of total expenditure on health	2001	17.9	World Health Report 2004, WHO
Total adult illiteracy rate	2000	42.8	UNESCO
Adult male illiteracy rate	2000	31.6	UNESCO
Adult female illiteracy rate	2000	54.6	UNESCO
Gross primary school enrolment ratio, male	2000/2001	not available	UNESCO
Gross primary school enrolment ratio, female	1996	not available	UNESCO
Gross secondary school enrolment ratio, male	1996	not available	UNESCO
Gross secondary school enrolment ratio, female	1996	not available	UNESCO

Contact address

UNAIDS/WHO Working Group on Global
HIV/AIDS and STI Surveillance
20, Avenue Appia
CH - 1211 Geneva 27
Switzerland
Fax: +41-22-791-4834

email: HIV-AIDS@who.int or estimates@unaids.org

website: <http://www.who.int/hiv>

<http://www.unaids.org>

Extracts of the information contained in these fact sheets may be reviewed, reproduced or translated for research or private study but not for sale or for use in conjunction with commercial purposes. Any use of information in these fact sheets should be accompanied by the following acknowledgment "WHO/UNAIDS epidemiological fact sheets on HIV/AIDS and Sexually Transmitted Infections, 2004 Update".

HIV prevalence in different populations

This section contains information about HIV prevalence in different populations. The data reported in the tables below are mainly based on the HIV database maintained by the United States Bureau of the Census where data from different sources, including national reports, scientific publications and international conferences are compiled. To provide a simple overview of the current situation and trends over time, summary data are given by population group, geographical area (Major Urban Areas versus Outside Major Urban Areas), and year of survey. Studies conducted in the same year are aggregated and the median prevalence rates (in percentages) are given for each of the categories. The maximum and minimum prevalence rates observed, as well as the total number of surveys/sentinel sites, are provided with the median, to give an overview of the diversity of HIV-prevalence results in a given population within the country. Data by sentinel site or specific study from which the medians were calculated are printed at the end of this fact sheet.

The differentiation between the two geographical areas Major Urban Areas and Outside Major Urban Areas is not based on strict criteria, such as the number of inhabitants. For most countries, Major Urban Areas were considered to be the capital city and - where applicable - other metropolitan areas with similar socio-economic patterns. The term Outside Major Urban Areas considers that most sentinel sites are not located in strictly rural areas, even if they are located in somewhat rural districts.

HIV sentinel surveillance*


Group	Area		1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003			
Pregnant women	Major urban areas	N-Sites		2.00	2.00	3.00	2.00	4.00	5.00	5.00	4.00	4.00	4.00	8.00	7.00	9.00	11.00	5.00				
		Minimum		0	0	0	0	0	0	0	0	0	0	0.50	0.25	0	0	0	0			
		Median		0	0	0	0	0.26	0	0.92	1.61	2.02	1.29	1.63	2.00	0.50	0.25	0.25				
		Maximum		0	0	0	0	1.28	0.80	2.50	2.25	2.38	3.00	3.75	3.25	6.50	7.75	1.25				
	Outside major urban areas	N-Sites	1.00	1.00	4.00	5.00	11.00	5.00	11.00	8.00	11.00	9.00	6.00	90.00	95.00	104.00	167.00	182.00				
		Minimum	0.05	0.10	0	0	0	0.08	0	0	0	0.25	0	0	0	0	0	0	0			
		Median	0.05	0.10	0	0.02	0.50	0.79	0	0.72	0.38	0.68	1.29	0.25	0.25	0.40	0.25	0.25				
		Maximum	0.05	0.10	0.04	0.38	2.44	3.80	1.91	1.13	4.25	3.00	3.62	28.24	6.50	4.25	5.25	8.75				
	Sex workers	Major urban areas	N-Sites	1.00	2.00	1.00	1.00	3.00	3.00	3.00	1.00	3.00	2.00	1.00	2.00	1.00	2.00	1.00				
			Minimum	0.89	4.45	13.20	21.11	0	0.72	1.41	1.58	1.20	11.27	11.93	5.30	64.40	58.67	52.26				
			Median	0.89	6.61	13.20	21.11	0.53	0.87	1.86	1.58	3.57	31.14	11.93	32.05	64.40	58.67	52.26				
			Maximum	0.89	8.76	13.20	21.11	33.93	7.58	50.99	1.58	4.80	51.00	11.93	58.80	64.40	58.67	52.26				
Outside major urban areas		N-Sites	5.00	3.00	5.00	4.00	5.00	7.00	6.00	3.00				1.00	2.00	3.00	4.00	4.00				
		Minimum	0	0	0	0	0	0	1.00	1.00				59.26	4.76	11.70	3.42	2.60				
		Median	0.49	1.42	4.94	16.83	19.00	23.71	23.25	29.00				59.26	6.53	13.10	25.85	29.80				
		Maximum	3.70	5.00	8.50	24.23	31.58	36.46	46.80	39.92				59.26	8.30	20.00	53.20	51.33				
Injecting drug users		Major urban areas	N-Sites					1.00	1.00	1.00		1.00	1.00				3.00	3.00	2.00			
			Minimum					1.92	1.51	0		1.65	3.52				5.00	2.40	1.46			
			Median					1.92	1.51	0		1.65	3.52				23.68	24.56	4.33			
			Maximum					1.92	1.51	0		1.65	3.52				26.67	41.38	7.20			
	Outside major urban areas	N-Sites	1.00	1.00	1.00	2.00	1.00	1.00	2.00	3.00	6.00	2.00	2.00	5.00	7.00	9.00	9.00	7.00	9.00			
		Minimum	0	0	8.57	39.12	44.75	67.20	1.50	25.41	3.90	55.71	67.10	0.96	1.20	1.41	1.39	0.40				
		Median	0	0	8.57	46.83	44.75	67.20	43.28	55.00	39.89	64.49	72.00	70.33	29.50	9.62	7.59	14.80				
		Maximum	0	0	8.57	54.54	44.75	67.20	85.06	85.64	61.05	73.27	76.90	76.10	68.40	75.60	59.60	51.60				
	STI patients	Major urban areas	N-Sites	2.00	2.00	3.00	8.00	7.00	9.00	11.00	7.00	3.00	4.00	8.00	6.00	6.00	8.00	11.00	5.00			
			Minimum	0	0.09	0.03	0	0.09	0.08	0.50	1.00	0.86	2.13	0	3.60	0.80	2.00	2.40	0.80			
			Median	0.52	4.88	0.27	1.86	5.30	3.20	3.68	25.55	5.65	15.95	2.78	20.00	6.35	3.84	6.40	3.60			
			Maximum	1.04	9.68	9.89	13.76	16.36	22.84	24.86	57.58	31.41	32.64	42.00	58.80	64.40	33.33	38.00	52.30			
Outside major urban areas		N-Sites	6.00	6.00	6.00	6.00	9.00	6.00	11.00	24.00	26.00	18.00	7.00	62.00	70.00	95.00	119.00	140.00				
		Minimum	0	0	0.94	0.60	1.33	2.60	0	0	0	0	4.85	0	0	0	0	0	0			
		Median	0.16	0.72	1.48	2.14	2.76	4.20	2.56	5.28	6.08	6.14	19.20	3.53	2.44	2.66	1.25	2.33				
		Maximum	0.80	1.42	3.02	4.80	8.84	13.75	6.19	20.50	21.66	19.31	22.40	50.25	44.80	48.00	41.60	39.20				
Men having sex with men		Major urban areas	N-Sites						2.00								2.00	2.00	1.00			
			Minimum						10.32								14.00	17.21	23.60			
			Median						13.11								18.97	20.41	23.60			
			Maximum						15.90								23.94	23.60	23.60			
	Outside major urban areas	N-Sites		2.00													1.00	1.00	2.00			
		Minimum		0													4.00	2.40	1.20			
		Median		0.19													4.00	2.40	2.40			
		Maximum		0.37													4.00	2.40	3.60			
	Tuberculosis patients	Major urban areas	N-Sites	1.00	1.00	3.00	2.00	3.00	4.00	2.00	2.00	2.00	3.00	3.00	1.00	1.00						
			Minimum	0	2.40	0	3.90	0.35	0	0	1.00	0.50	1.50	2.86	30.00	33.00						
			Median	0	2.40	2.30	3.93	7.07	4.59	5.30	6.00	5.60	3.20	3.40	30.00	33.00						
			Maximum	0	2.40	3.79	3.96	7.90	10.60	10.60	11.00	10.70	16.80	23.00	30.00	33.00						
Outside major urban areas		N-Sites					1.00	7.00	4.00	4.00	6.00	4.00	3.00	6.00	4.00			1.00				
		Minimum					3.20	0.71	0	3.20	2.00	3.60	4.39	6.00	2.80			4.50				
		Median					3.20	1.10	4.13	7.14	4.22	13.79	21.43	9.48	13.00			4.50				
		Maximum					3.20	4.46	11.50	14.30	12.10	18.50	21.50	17.60	20.06			4.50				

*Detailed data by site can be found in the Annex.


Maps & charts

Mapping the geographical distribution of HIV prevalence among different population groups may assist in interpreting both the national coverage of the HIV surveillance system as well in explaining differences in levels of prevalence. The UNAIDS/WHO Working Group on Global HIV/AIDS and STI Surveillance, in collaboration with the WHO Public Health Mapping Team, Communicable Diseases, is producing maps showing the location and HIV prevalence in relation to population density, major urban areas and communication routes. For generalized epidemics, these maps show the location of prevalence of antenatal surveillance sites.

Trends in antenatal sentinel surveillance for higher prevalence countries, or in prevalence among selected populations for countries with concentrated epidemics, are a new addition. These are presented for those countries where sufficient data exist.



Median HIV prevalence in ANC population in Andhra Pradesh, Karnataka, Maharashtra and Tamil Nadu, India. 1998-2003*



Median prevalence and ranges are shown in areas with more than one sentinel site.

The boundaries and names shown and the designations used on the map do not imply the expression of any opinion whatsoever on the part of the World Health Organization concerning the legal status of any country, territory, city or area or of its authorities, or concerning the delimitation of its frontiers or boundaries. Dotted lines on maps represent approximate border lines for which there may not yet be full agreement. WHO 2004, all rights reserved.

Reported AIDS cases

Following WHO and UNAIDS recommendations, AIDS case reporting is carried out in most countries. Data from individual AIDS cases are aggregated at the national level and sent to WHO. However, case reports come from surveillance systems of varying quality. Reporting rates vary substantially from country to country and low reporting rates are common in developing countries due to weaknesses in the health care and epidemiological systems. In addition, countries use different AIDS case definitions. A main disadvantage of AIDS case reporting is that it only provides information on transmission patterns and levels of infection approximately 5-10 years in the past, limiting its usefulness for monitoring recent HIV infections.

Despite these caveats, AIDS case reporting remains an important advocacy tool and is useful in estimating the burden of HIV-related morbidity as well as for short-term planning of health care services. AIDS case reports also provide information on the demographic and geographic characteristics of the affected population and on the relative importance of the various exposure risks. In some situations, AIDS reports can be used to estimate earlier HIV infection patterns using back-calculation. AIDS case reports and AIDS deaths have been dramatically reduced in industrialized countries with the introduction of Anti-Retroviral Therapy (ART).

1979	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998
0	0	0	0	0	0	0	0	0	0	0	57	45	140	252	523	1091	888	2108	1148
1999	2000	2001	2002	2003	Total		UNK	Date of last report											
					8438		2186	8/31/1999											

Curable sexually transmitted infections (STIs)

The predominant mode of transmission of both HIV and other STIs is sexual intercourse. Measures for preventing sexual transmission of HIV and STIs are the same, as are the target audiences for interventions. In addition, strong evidence supports several biological mechanisms through which STIs facilitate HIV transmission by increasing both HIV infectiousness and HIV susceptibility. Thus, detection and treatment of individuals with STIs is an important part of an HIV control strategy. In summary, if the incidence/prevalence of STIs is high in a country, then there is the possibility of high rates of sexual transmission of HIV. Monitoring trends in STIs provides valuable insight into the likelihood of the importance of sexual transmission of HIV within a country, and is part of second generation surveillance. These trends also assist in assessing the impact of behavioural interventions, such as delaying sexual debut, reducing the number of sex partners and promoting condom use.

Clinical services offering STI care are an important access point for people at high risk for both STIs and HIV. Identifying people with STIs allows for not only the benefit of treating the STI, but for prevention education, HIV testing, identifying HIV-infected persons in need of care, and partner notification for STIs or HIV infection. Consequently, monitoring different components of STI prevention and control can also provide information on HIV prevention and control activities within a country.

STI syndromes

Reported cases	1996	1997	1998	1999	2000	2001	2002	2003	Incidence 2003
----------------	------	------	------	------	------	------	------	------	----------------

Comments:

Source:

Syphilis prevalence, women

Percent of blood samples taken from pregnant women aged 15-49 that test positive for syphilis - positive reaginic and treponemal test-during routine screening at selected antenatal clinics.

Year	Area	Rate	Range
------	------	------	-------

Comments:

Source:

Estimated prevalence of curable STIs among female sex workers

- Chlamydia

Year	Area	Rate	Range
------	------	------	-------

Comments:

Source:

- Gonorrhoea

Year	Area	Rate	Range
------	------	------	-------

Comments:

Source:

Estimated prevalence of curable STIs among female sex workers (continued)**- Syphilis**

Year	Area	Rate	Range
------	------	------	-------

Comments:

Source:

- Trichomoniasis

Year	Area	Rate	Range
------	------	------	-------

Comments:

Source:

Health service and care indicators

HIV prevention strategies depend on the twin efforts of care and support for those living with HIV or AIDS, and targeted prevention for all people at risk or vulnerable to the infection. It is difficult to capture such a large range of activities with one or just a few indicators. However, a set of well-established health care indicators may help to identify general strengths and weaknesses of health systems. Specific indicators, such as access to testing and blood screening for HIV, help to measure the capacity of health services to respond to HIV/AIDS - related issues.

Access to health care

Indicators	Year	Estimate	Source
% of population with access to health services - total			
% of population with access to health services - urban			
% of population with access to health services - rural			
Contraceptive prevalence rate (%)	1999	48.2	UNICEF/UNPOP
Percentage of contraceptive users using condoms			
% of births attended by skilled health personnel	2000	42.3	WHO
% of 1-yr-old children fully immunized - DPT	2002	70	WHO/UNICEF
% of 1-yr-old children fully immunized - Measles	2001	56	WHO/UNICEF
% of ANC clinics where HIV testing is available			

Number of adults (15-49) with advanced HIV infection receiving ARV therapy as of June 2004

Adults on treatment

Number: 21000

Source: WHO

Estimated number of adults (15-49) in need of treatment in 2003

Adults needing treatment

Number: 710,000

Source: WHO/UNAIDS

Coverage of HIV testing and counselling

Number of public and NGO services providing testing and counselling services.

Year	Area	N=
------	------	----

Comments:

Source:

Knowledge and behaviour

In most countries the HIV epidemic is driven by behaviours (e.g.: multiple sexual partners, injecting drug use) that expose individuals to the risk of infection. Information on knowledge and on the level and intensity of risk behaviour related to HIV/AIDS is essential in identifying populations most at risk for HIV infection and in better understanding the dynamics of the epidemic. It is also critical information in assessing changes over time as a result of prevention efforts. One of the main goals of the 2nd generation HIV surveillance systems is the promotion of a standard set of indicators defined in the National Guide (Source: National AIDS Programmes, A Guide to Monitoring and Evaluation, UNAIDS/00.17) and regular behavioural surveys in order to monitor trends in behaviours and to target interventions.

The indicators on knowledge and misconceptions are an important prerequisite for prevention programmes to focus on increasing people's knowledge about sexual transmission, and, to overcome the misconceptions that act as a disincentive to behaviour change. Indicators on sexual behaviour and the promotion of safer sexual behaviour are at the core of AIDS programmes, particularly with young people who are not yet sexually active or are embarking on their sexual lives, and who are more amenable to behavioural change than adults. Finally, higher risk male-male sex reports on unprotected anal intercourse, the highest risk behaviour for HIV among men who have sex with men.

Knowledge of HIV prevention methods

Prevention indicator: Percentage of young people 15-24 who both correctly identify two ways of preventing the sexual transmission of HIV and who reject three misconceptions about HIV transmission.

Year	Male	Female
2000	17	21

Comments:

Source: BSS

Reported condom use at last higher risk sex (young people 15-24)

Prevention indicator: Proportion of young people reporting the use of a condom during sex with a non-regular partner.

Year	Male	Female
2001	59	51

Comments:

Source: BSS

Age-mixing in sexual partnerships among young women

The proportion of young women who have had sex in the last 12 months with a partner who is 10 or more years older than themselves.

Year	Area	Age group	Male	Female	All
------	------	-----------	------	--------	-----

Comments:

Source:

Reported non-regular sexual partnerships

Prevention indicator: Proportion of young people 15-24 having at least one sex partner other than a regular partner in the last 12 months.

Year	Male	Female
2001	12	2

Comments:

Source: BSS

Knowledge and behaviour (continued)Ever used a condom

Percentage of people who ever used a condom.

Year	Area	Age group	Male	Female	All
------	------	-----------	------	--------	-----

Comments:

Source:

Adolescent pregnancy

Percentage of teenagers 15-19 who are mothers or pregnant with their first child.

Year	Percentage
------	------------

Comments:

Source:

Age at first sexual experience

Proportion of 15-19 year olds who have had sex before age 15.

Year	Male	Female
------	------	--------

Comments:

Source:

Prevention indicators

Male and female condoms are the only technology available that can prevent sexual transmission of HIV and other STIs. Persons exposing themselves to the risk of sexual transmission of HIV should have consistent access to high quality condoms. AIDS Programs implement activities to increase both availability of and access to condoms. These activities should be monitored and have resources directed to problem areas. The indicator below highlights the availability of condoms. However, even if condoms are widely available, this does not mean that individuals can or do access them.

Condom availability nationwide

Total number of condoms available for distribution nationwide during the preceding 12 months, divided by the total population aged 15-49.

Year	N	Rate
------	---	------

Comments:

Source:

Prevention of mother-to-child transmission (MTCT) nationwide

Percentage of women who were counselled during antenatal care for their most recent pregnancy, accepted an offer of testing and received their test results, of all women who were pregnant at any time in the preceding two years.

Year	N	Rate
------	---	------

Comments:

Source:

Blood safety programs aim to ensure that the majority of blood units are screened for HIV and other infectious agents. This indicator gives an idea of the overall percentage of blood units that have been screened to high enough standards that they can confidently be declared free of HIV.

Screening of blood transfusions nationwide

Percentage of blood units transfused in the last 12 months that have been adequately screened for HIV according to national or WHO guidelines.

Year	N	Rate
------	---	------

Comments:

Source:

Sources

Data presented in this Epidemiological Fact Sheet come from several sources, including global, regional and country reports, published documents and articles, posters and presentations at international conferences, and estimates produced by UNAIDS, WHO and other United Nations agencies. This section contains a list of the more relevant sources used for the preparation of the Fact Sheet. Where available, it also lists selected national Web sites where additional information on HIV/AIDS and STI are presented and regularly updated. However, UNAIDS and WHO do not warrant that the information in these sites is complete and correct and shall not be liable whatsoever for any damages incurred as a result of their use.

Aher, A. 2002 Intervention amongst MSMs at Surface Railway Stations in Mumbai Metro XIV International AIDS Conference, Barcelona, Spain, 7/7-12, Abstract ThPeD7718.

Ambwani, P. N., I. S. Gilada, J. Karkare, et al. 1998 Prevention of Perinatal Transmission: IHO-Wadia Model 12th World AIDS Conference, Geneva, 6/28-7/3, Abstract 23309.

Babu, P. G., T. Ishida, V. Nerurkar, et al. 1994 Epidemiology of Retroviral Infections in South India Tenth International Conference on AIDS, Yokohama, Japan, 8/7-12, Poster P.C.0082.

Babu, P. G., T. Ishida, V. Nerurkar, et al. 1994 Epidemiology of Retroviral Infections in South India Tenth International Conference on AIDS, Yokohama, Japan, 8/7-12, Poster P.C.0082.

Behra, R., N. M. Mehta, A. Purohit, et al. 2000 Study of Incidence of HIV Seropositivity at the Blood Bank in Jodhpur, Rajassthan, India XIII International AIDS Conference, Durban, South Africa, 7/9-14, Abstract MoPeC2422.

Bhave, G., S. Desai, V. Parkar 1996 Trends of HIV-1 and HIV-2 Infection in STD Patients and Pattern of Transmission to Their Spouses in Bombay, India XI International Conference on AIDS, Vancouver, 7/7-14, Poster, Mo.C.1496.

Bhurushundi, M., R. Munje, A. Saraf, et al. 2000 HIV Seropositivity in Tuberculosis XIII International AIDS Conference, Durban, South Africa, 7/9-14, Abstract WePeC4412.

Bhuyan, K. K., G. A. Dallabetta, S. J. Mills 2001 Comparison of Prevalence of STI/HIV and Their Behavioral Correlates among Female Sex Workers from Three States of India 6th International Congress on AIDS in Asia and the Pacific, Melbourne, Australia, 10/5-10, Poster Sa1672.

Chakrabarty, M. S., P. N. Dey, S. Paul, et al. 1994 Seroepidemiology of HIV Infection in Calcutta Tenth International Conference on AIDS, Yokohama, Japan, 8/7-12, Poster P.C.0083.

Datta, M. 1995 Tuberculosis Related to HIV/AIDS 3rd International Conference on AIDS in Asia and the Pacific, Chiang Mai, Thailand, 9/17-21, Session WS10.

Dey, S. K. 1999 Spectrum of HIV Infection among STD Clinic Attenders: Experiences of Different Perspectives from Rural North Bengal and Urban .. 5th International Congress on AIDS in Asia and the Pacific, Kuala Lumpur, Malaysia, 10/20-27, Abstract PTAB034.

Dey, S. K., N. K. P. Pal, N. B. Bhattacharjee, et al. 1998 Spectrum of HIV Infection in Tuberculosis (TB) - Urban and Rural Experiences of Different Perspectives, West Bengal, India 12th World AIDS Conference, Geneva, 6/28 - 7/3, Abstract 22152.

Dey, S. K., N. K. P. Pal, N. B. Bhattacharjee, et al. 1998 Changing Pattern of HIV Infection in Some Risk Groups: 1987-1996, West Bengal, India 12th World AIDS Conference, Geneva, 6/28 - 7/3, Poster 43491.

Dey, S. K., N. Pal, U. Ganguly, et al. 1999 Measuring the Impact of Interventions in HIV Epidemic - Experience from a Rural Belt of North Bengal and Urban Calcutta, India 5th International Congress on AIDS in Asia and the Pacific, Kuala Lumpur, Malaysia, 10/20-27, Abstract PSCD111.

Ekstrand, M., C. Lindan, G. Bhave, et al. 1998 Male STD Patients in Mumbai, India are in Urgent Need of Culturally Specific and Feasible AIDS Prevention Programs 12th World AIDS Conference, Geneva, 6/28 - 7/3, Abstract 14209.

Gadkari, A. D., et al. 1995 HIV Seroprevalence in STD and TB Clinics TB & HIV, no. 8, p. 28.

Gandhi, N. R., K. Mayer, A. Patel, et al. 2001 HIV Screening in Active Tuberculosis Patients in Rural Gujarat, India International Journal of STD and AIDS, vol. 12, no. 6, pp. 415-416.

Gilada, I., R. Mahajan, S. Hira 1994 HIV Infection in Pregnant Women in Bombay Tenth International Conference on AIDS, Yokohama, Japan, 8/7-12, Poster P.C.0081.

Gopalakrishnan, P. B., S. Padmarajan, M. Johnson, et al. 1992 HIV the 3rd Generation STD Problem 2nd International Congress on AIDS in Asia and Pacific, New Delhi, India, 11/8-12, Poster A403.

Hussain, M., N. Islam, Z. Hossain, et al. 1999 Study on Prevalence of HBV, HCV and HIV Markers among Blood Donors and Pregnant Women in Bangladesh Department of Transfusion Medicine, IPGMR, Dhaka, Bangladesh, November, report.

Ingle, K., Gaikwad, G. Jadhao, et al. 2001 HIV Seropositivity in Tuberculosis in Rural District, India 6th International Congress on AIDS in Asia and the Pacific, Melbourne, Australia, 10/5-10, Abstract 0452.

Jacob, M., T. J. John, G. Soshamma, et al. 1995 Increasing Prevalence of Human Immunodeficiency Virus Infection among Patients Attending a Clinic for Sexually Transmitted . . . Indian Journal of Medical Research, vol. 101, pp. 6-9.

Jagavkar, C., P. Dalaa, A. Chowdhary 1995 GUD and HIV Infection in STD Centennial Surveillance in Bombay IUVDT World STD/AIDS Congress, Singapore, 3/19-23, Free Paper 11.

Jagtap, M. 1995 HIV Epidemic in Maharashtra State, India 3rd International Conference on AIDS in Asia and the Pacific, Chiang Mai, Thailand, 9/17-21, Abstract PB115.

Jerajani, H., M. S. Setia, R. S. Dhurat 2000 A Study of the Trends of Clinic Attendance in an STD Clinic XIII International AIDS Conference, Durban, South Africa, 7/9-14, Poster WePeC4363.

John, T. J., N. Bhushan, P. G. Babu, et al. 1993 Prevalence of HIV Infection in Pregnant Women in Vellore Region Indian Journal of Medical Research, vol. 97, pp. 227-230.

John, T. J., P. G. Babu, H. Jayakumari, et al. 1987 Prevalence of HIV Infection in Risk Groups in Tamil Nadu, India Lancet, Jan. 17, vol. 1, no. 8525, pp. 160-161.

Johns Hopkins University 1986 AIDS -- A Public Health Crisis Population Information Program, Population Reports, Issues in World Health, July-Aug., Series L, no. 6, pp. 194-228.

Kamat, H. A., D. D. Banker 1993 Human Immunodeficiency Virus-1 Infection among Patients with Sexually Transmitted Diseases in Bombay The National Medical Journal of India, vol. 6, no. 1, pp. 11-13.

Kant, S., M. Shaukat, P. Salil 2000 Validation of an Assumption Used for Estimating HIV Prevalence in India XIII International AIDS Conference, Durban, South Africa, 7/9-14, Poster MoPeC2479.

Khan, M. A. 1998 Behavioural Aspects of HIV Infection amongst the Sex Workers of Agra 12th World AIDS Conference, Geneva, 6/28 - 7/3, Poster 23549.

Krishnamurthy, P. 1999 Community Prevalence of STD - in Tamil Nadu, India Presented at Monitoring the AIDS Pandemic (MAP) in Asia Symposium, Network Consultative Meeting, 10/19-21, Kuala Lumpur, Malaysia.

Kura, M., M. Kohli, R. D. Kulkarni, et al. 1996 Pattern of STD in Bombay XI International Conference on AIDS, Vancouver, 7/7-14, Abstract Pub.C.1271.

Lakshmi, N., A. G. Kumar 1991 HIV Infection in Tirupati, India Genitourinary Medicine, vol. 67, pp. 427-428.

Lal, S., et al. 1991 AIDS Control Programme of India Government of India, Nirman Bhawan, New Delhi, India, Unpublished report.

Lal, S., L. Khodakevich, P. Salil 1994 HIV Infection in India - Trends Analysis Tenth International Conference on AIDS, Yokohama, Japan, 8/7-12, Session 039C.

Li, P. C., E. K. Yeoh 1992 Current Epidemiological Trends of HIV Infection in Asia AIDS Clinical Review, pp. 1-23.

Lindan, C., G. Bhave, P. Gupte, et al. 1997 High Prevalence of HIV among Male STD Patients in Bombay Heralds Spread to the General Population 4th International Congress on AIDS in Asia and the Pacific, Manila, Philippines, 10/25-29, Abstract BP120.

Marques, L., A. Purohit 1999 Counseling Plays an Important Role in HIV Prevention among CSWs 5th International Congress on AIDS in Asia and the Pacific, Kuala Lumpur, Malaysia, 10/20-27, Abstract PMCD086.

Mathur, D., P. Durlabhji, N. M. Singhvi, et al. 1997 HIV Sentinel Surveillance in STD and Antenatal Clinic Attenders 4th International Congress on AIDS in Asia and the Pacific, Manila, Philippines, 10/25-29, Abstract AP115.

Mathur, R., P. V. Roa, Y. N. Mathur 1988 Prevalence of HIV Antibody in High Risk Groups, Jabalpur, Central India Virus Information Exchange Newsletter, vol. 5, no. 2, p. 61.

Mathur, S. C., S. Pathak, A. Mathur 2000 HIV Sentinel Surveillance in Rajasthan XIII International AIDS Conference, Durban, South Africa, 7/9-14, Abstract MoPeC2457.

Mehendale, S., J. J. Rodrigues, R. Gangakhedkar, et al. 1994 STDs and HIV Infection in CSWs of Pune, India Tenth International Conference on AIDS, Yokohama, Japan, 8/7-12, Abstract P.C.0351.

Mehta, N. M., A. Purohit, J. Chakraborty, et al. 1999 Update of the Study of Incidence of HIV Seropositivity at the Blood Bank in Jodhpur, India 5th International Congress on AIDS in Asia and the Pacific, Kuala Lumpur, Malaysia, 10/20-27, Abstract PTCD009.

Merchant, R. H., K. Damanika, I. S. Gilada, et al. 2001 Strategy for Preventing Vertical Transmissions of HIV: Bombay Experience Indian Pediatrics, vol. 38, pp. 132-138.

Mohanty, K. C., R. M. Sundrani 2000 Prevalence of Tuberculosis & HIV in Indoor Patients XIII International AIDS Conference, Durban, South Africa, 7/9-14, Abstract WePeC4419.

Moses, S. 2001 Enabling a "Healthy Highway" for Truckers in Visakhapatnam, India 6th International Congress on AIDS in Asia and the Pacific, Melbourne, Australia, 10/5-10, Poster Su2213.

Murugasampillay, S. 1993 HIV and AIDS Surveillance at State Level in India a Public Health Tool for AIDS and STD Control National AIDS Control Organisation - India and Global Programme on AIDS, World Health Organisation, 12th August - 8th December, draft report.

Narain, J. P., A. Jha, S. Lal. et al. 1994 Risk Factors for HIV Transmission in India AIDS, vol. 8, suppl. 2, pp. S77-S82.

National AIDS Control Programme 1997 HIV Sentinel Surveillance Report National AIDS Control Programme, India, unpublished document.

National AIDS Control Organisation 1998 Country Scenario 1997-98 National AIDS Control Organisation, Ministry of Health and Family Welfare, Government of India, pp. 15-31.

National AIDS Control Organisation 1998 HIV Sentinel Surveillance Report: August - October 1999 National AIDS Control Organisation, India, unpublished tables.

National AIDS Control Programme 2001 Current Status and Trend of HIV/AIDS Epidemic in India: 2001 National AIDS Control Organization, Ministry of Health and Family Welfare, Government of India, report.

National AIDS Control Programme 2001 HIV Sentinel Surveillance Report National AIDS Control Organisation, India, unpublished tables.

National AIDS Control Organization 2003 HIV Sentinel Surveillance Report Population and Sentinel Site - Wise Report: 1998-2002 National AIDS Control Organization, India, May, unpublished tables.

National STD Control Programme 1995 National STD Control Programme Ministry of Health and Family Welfare.

Oberai, C., N. Daithankar, M. Kohli, et al. 1995 Epidemiologic Features of HIV-Infection in a Referral Clinic in Bombay IUVDT World STD/AIDS Congress, Singapore, 3/19-23, Free Paper 9.

Pal, N. K., A. Das, P. K. Halder, et al. 1995 HIV and Syphilis in CSWs Clients and IV Drug Abusers in Calcutta 3rd International Conference on AIDS in Asia and the Pacific, Chiang Mai, Thailand, 9/17-21, Poster PB123.

Pal, S. C., S. Sarkar, T. N. Naik, et al. 1990 Explosive Epidemic of HIV Infection in North Eastern States of India, Manipur and Nagaland CARC Calling, vol. 3, no. 3, pp. 2-6.

Palaniappan, K. 1995 Trend of HIV among STD Patients, Pregnant Women and Truckers through Unlinked Anonymous Screening in India 3rd International Conference on AIDS in Asia and the Pacific, Chiang Mai, Thailand, 9/17-21, Poster PB120.

Panda, S., G. Kame, M. Pamei, et al. 1994 Clinical Features of HIV Infection in Drug Users of Manipur National Medical Journal of India, vol. 7, no. 6, pp. 267-269.

Pankajalakshmi, V. V., A. Uma, R. Sethuraman, et al. 1992 HIV Seropositivity among STD Patients 2nd International Congress on AIDS in Asia and Pacific, New Delhi, India, 11/8-12, Poster B704.

Paranjape, R. S., S. P. Tripathy, P. A. Menon, et al. 1997 Increasing Trend of HIV Seroprevalence among Pulmonary Tuberculosis Patients in Pune, India Indian Journal of Medical Research, vol. 106, pp. 207-211.

Pedhambkar, R. B., B. S. Pedhambkar, and M. M. Kura 2001 Study of Risk Factors Associated with HIV Seropositivity in STD Patients at Mumbai, India Sexually Transmitted Infections, vol. 77, no. 5, pp. 388-389.

Porterfield, D. S., A. D. Nageswari, R. Paramasivam, et al. 1992 HIV Infection, Sexual Habits and Tuberculosis in South India 2nd International Congress on AIDS in Asia and Pacific, New Delhi, India, 11/8-12, Abstract B331.

Purohit, A., N. M. Mehta, R. Behra, et al. 1998 Incidence of HIV Seropositivity at the Blood Bank in Jodhpur, India 12th World AIDS Conference, Geneva, 6/28-7/3, Poster 23259.

Rajan, R. 1992 Documentation - Need within Different User Category 2nd International Congress on AIDS in Asia and Pacific, New Delhi, India, 11/8-12, Poster D205.

Raviglione, M. C., F. Luelmo 1996 Update on the Global Epidemiology of Tuberculosis Current Issues in Public Health, vol. 2, pp. 192-197.

Ray, K., V. Ramesh, S. N. Karmakar, et al. 1996 Increasing Trend of HIV Seropositivity in a Sexually Transmitted Diseases Centre and Epidemiology of HIV Seropositive ... International Journal of STD and AIDS, vol. 7, no. 1, pp. 48-50.

Rose, A., H. Srinivasa, R. S. Macaden, et al. 1992 Anonymous HIV Screening of Pregnant Women, Women with Bad Obstetric History and Patients from Psychiatry 2nd International Congress on AIDS in Asia and Pacific, New Delhi, India, 11/8-12, Abstract A602.

Salunke, S., M. R. Jagtap, S. Hira, et al. 1997 Rapid Rise in HIV Prevalence among Women Attending Government STD Clinic in Mumbai (Bombay), India International Journal of STD and AIDS, vol. 8, no. 4, p. 280.

Samani, P. H., S. Sankaranarayanan, A. Gogate 2002 Analytical Study of Clinical Case Profile of Opportunistic Infections (OIs) in People Living with HIV/AIDS (PLWH/A) in Mumbai XIV International AIDS Conference, Barcelona, Spain, 7/7-12, Abstract ThPeF8215.

Samaram, G. 2001 A Study of HIV/AIDS in Vijayawada - AP India 6th International Congress on AIDS in Asia and the Pacific, Melbourne, Australia, 10/5-10, Poster.

Sankari, S., S. Solomon, et al. 1991 Trends of HIV Infections in Antenatal/Infertility Clinic - An Ominous Sign: VII International Conference on AIDS, Florence, Italy, 6/16-21, Poster W.C.3236.

Sarkar, S., N. Das, S. Panda, et al. 1993 Rapid Spread of HIV among Injecting Drug Users in North-Eastern States of India Bulletin on Narcotics, vol. XLV, no. 1, pp. 91-105.

Sato, P. 1992 HIV Sero-Survey Data, India Presented/discussed during Calcutta Workshop, October 28, document.

Saxena, D. M., J. K. Kosambiya 1992 HIV Seropositivity in Sex Workers of Surat 2nd International Congress on AIDS in Asia and Pacific, New Delhi, India, 11/8-12, Poster B338.

Sengupta, S., C. Priyamvada 1997 HIV Surveillance in Poor Women of the General Population Vis-a-Vis STD Prevalence 4th International Congress on AIDS in Asia and the Pacific, Manila, Philippines, 10/25-29, Poster AP061.

Setia, M., H. R. Jerajani, A. R. Kavi, et al. 2001 Male STI Patients in Mumbai, India are in Urgent Need of HIV Prevention Programs 6th International Congress on AIDS in Asia and the Pacific, Melbourne, Australia, 10/5-10, Poster Su1350.

Shaukat, M. 1999 Current Status and Trend of HIV/AIDS Epidemic in India Presented at Monitoring the AIDS Pandemic (MAP) in Asia Symposium, Network Consultative Meeting, 10/19-21, Kuala Lumpur, Malaysia.

Shepherd, M. E., S. M. Mehendale, R. S. Paranjape, et al. 2002 Stable HIV Incidence Over an 8-Year Period among Male Patients Attending STD Clinics in Pune India XIV International AIDS Conference, Barcelona, Spain, 7/7-12, Poster TuPeC4899.

- Simoes, E. A. F., P. G. Babu, H. M. Jeyakumari, et al. 1993 The Initial Detection of Human Immunodeficiency Virus 1 and its Subsequent Spread in Prostitutes in Tamil Nadu, India Journal of Acquired Immune Deficiency Syndromes, vol. 6, no. 9, pp. 1030-1034.
- Singh, B. 1988 Screening of HIV Antibody among High Risk Groups in Manipur IV International Conference on AIDS, Stockholm, 6/15-16, Abstract 5509.
- Singh, N. B., Y. I. Singh, H. L. Singh 1991 Epidemic of HIV Infection among Intravenous Drug Users in Manipur, India Virus Information Exchange Newsletter, vol. 8, no. 1, p. 20.
- Solomon, S., S. Anuradha, M. Ganapathy, et al. 1994 Sentinel Surveillance of HIV-1 Infection in Tamilnadu, India International Journal of STD and AIDS, vol. 5, pp. 445-446.
- Solomon, S., Sundararaman, G. Babu, et al. 1988 Quantitative Estimation of HIV Antibodies in Prostitutes IV International Conference on AIDS, Stockholm, 6/15-16, Poster 5512.
- Tripathy, S. P., D. Joshi, P. Menon, et al. 1996 Seroprevalence of HIV-1 Infection in Tuberculosis Patients at Pune, India XI International Conference on AIDS, Vancouver, 7/7-14, Abstract Pub.B.1091.
- Tripathy, S. P., D. R. Joshi, P. Menon, et al. 1997 Sentinel Surveillance for HIV Infection in Tuberculosis Patients in Pune, India 4th International Congress on AIDS in Asia and the Pacific, Manila, Philippines, 10/25-29, Abstract Do041.
- Tripathy, S., K. Banerjee, J. Rodrigues, et al. 1993 Increasing HIV Infection in Western India IX International Conference on AIDS, Berlin, 6/6-11, Poster PO-C08-2764.
- U.S. Department of State 2000 HIV/AIDS in Northeast India Unclassified cable, December, Calcutta 0610.
- Verenkar, M., S. Rodrigues, M. J. Pinto, et al. 1992 HIV, Hepatitis B and Syphilis among Sex Workers of Goa 2nd International Congress on AIDS in Asia and Pacific, New Delhi, India, 11/8-12, Poster A128.

Websites: **National AIDS Control Organization (NACO):** <http://www.naco.nic.in/>
 Indian Health Organization (IHO): <http://www.wwindia.com/iho/>
 Ministry of Health and Family Welfare: <http://mohfw.nic.in/>

Annex: HIV surveillance by site

Group	Area	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003		
Pregnant women	Outside major urban areas	District hospital, Leh											0	0		0				
		District hospital, Mandasaur												0	0.50	0.50	1.00			
		District hospital, Murshidabad													0	0.63				
		District hospital, Nayagaon												0	0	0	0			
		District hospital, Puruliya																0		
		District hospital, Raichur																2.75		
		District hospital, Raigarh																0		
		District hospital, Rajnandgaon																	0.54	
		District hospital, Ratlam													0	0	0.25	0.25		
		District hospital, Rewa													0	0.25	0	0.75		
		District hospital, Sagar																	0.25	
		District hospital, Shahdol													0	0.25	0	0		
		District hospital, Shivpuri																	0	
		District hospital, Solan															1.23	0		
		District hospital, Srigangagar																	0	
		District hospital, Surendranagar																	0.50	
		District hospital, Tamenglong																	1.33	
		District hospital, Udipi															1.25	0.75		
		Diu																		0.22
		Duncan Hospital, Raxaul													0	0	0.25	0.25		
		Dungarpur																		0.75
		Eluru																		2.25
		Etawah																		3.00
		Faridkot																		0.25
		G. D. Hospital, Shillong													0.25	0	0	0.25		
		Gandhi MC Hospital, Hyderabad															2.00	0.50		
		Gangtok																		0
		GB hospital, Agartala																0.25		
		GB Pant Hospital, Port Blair													0	0	0	0		
		General Hospital, Hissar															0	0.25		
		General Hospital, Jammu															0.25	0.75		
		General Hospital, Kollegal, Mysore														0.25	1.61	1.00		
		General hospital, Palwai															0	0.50		
		General Hospital, Sector 16, Chandni													0.47	0.75	0	0		
		GH, Coimbatore																		0.25
		Gorakhpur																		0.25
		Government hospital, Akola													1.50	0.25	0.75	0.50		
		Government hospital, Along															0	0		
		Government hospital, Behrampur																		0.13
		Government hospital, Burla																		0.26
		Government hospital, Cuddalore																		1.00
		Government hospital, Dindigul																		0.50
		Government hospital, Itanagar													0.45	0				
		Government hospital, Karnal													0	0	0.25	0.50		
		Government hospital, Kolhapur													5.00	3.00	1.00	4.00		
		Government hospital, Minicoy													1.33	0	0	0		

Group	Area	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003		
Pregnant women	Outside major urban areas	Government hospital, Nasik											2.50	1.75	1.25	3.75				
		Government hospital, Pudukottai															0.75			
		Government hospital, Ratnagiri												0.50	0.75		1.01			
		Government hospital, Rourkela													0.25	0.30	0.26			
		Government hospital, Satara												2.25	3.75	4.25	2.25			
		Government hospital, Thodupuzha												0	0	0	0			
		Government hospital, Vellore																0.75		
		Government Medical College, Auranga													0	1.50	0.50	0.25	1.25	
		Government Medical College, Sholapu													1.00	1.50	2.25	1.50		
		Guntur																		4.00
		Haldwani																	0	0
		Hamirpur																		0.75
		Harda																		0
		Haridwar																		0.74
		Hassan																		0.75
		Himachal Pradesh state													0.40					
		Himmat Nagar																		0.75
		Hissar																		0
		Hoshiarpur																		0
		Hyderabad																		2.00
		IG Hospital, Kavaratti													0	0	0	0		
		IGMC, Nagpur															0.75	1.25		
		Imphal						1.00	1.00											0
		Institute of OB & GYN, Egmore, Chen													0.79	1.25	1.00	1.50		
		Institute of Prevention Medicine, H													2.50	3.25				
		Jagatsinghpur																		0
		Jagdulpur																		0.25
		Jaipur																		0.50
		Jammu																		0.25
		Jamshedpur																		0
		Jay Kay Lon Medical College, Kota													0	0.50	0	0		
		Jhalawar																		0.25
		Jind																		1.50
		JIPMER, Pondicherry														1.50	0.25			
		Jiribam																		0.25
		Jodhpur																		2.25
		Jorhat																		0
		Junagadh																		0.25
		Kakatia Medical College, Warangal															1.25	1.50		
		Kakinada																		3.00
		Kannur																		0
		Kanpur																		0
		Karimnagar																		1.50
		Karnal																		0
		Karnataka Medical College, Hubli													1.75	2.00				
		Katni																		0
		Kavaratti										0								0
		Kelong																		0
		KIMS, Hubli															1.75	1.00	1.75	
		Kinnaur																		0
		Kohima										0.50								1.14
		Kothagudem																		0.25
		Kottayam																		0.50
		Kurnool																		0.50
		Lal Ded Hospital, Srinagar													0	0	0	0		
		Lalitpur																		0.50
		Latur, Sholapur																		0.75
		Leh																		0
		Lucknow																		0
		Ludhiana																		0.49

Group	Area	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003		
Pregnant women	Outside major urban areas	Lungei																0.50		
		Machillipatnam																	2.25	
		Madurai																	1.00	
		Madurai Medical College, Madurai												1.00	0.75	0.50	1.25			
		Madurai Medical College, Micro. & V			0		0.15	0.28												
		Madurai Medical College, Coimbatore															0.50			
		Maharashtra state								0.80										
		Mahbubnagar																		0.25
		Mahila Hospital, Aligarh													0.23	0	0.25	0		
		Mahila Hospital, Bahraich																		0
		Mahila Hospital, Ballia																		0
		Mahila Hospital, Bijnore																		0
		Mahila Hospital, Etawah																		0.25
		Mahila Hospital, Gorakhpur													0	0	0	0.50		
		Mahila Hospital, Haldwani													0	0	0			
		Mahila Hospital, Mirzapur																		0
		Mahila Hospital, Nainital																		0
		Mahila Hospital, Partapgarh																		0.50
		Mahila Hospital, Pithoragarh																		0
		Mandi																		0
		Mandsaur																		0.25
		Maskati Hospital, Surat															0.75	0.25		
		Maternity hospital, Pondicherry													0.50	0.25	0.25			
		Maternity hospital, Pondicherry Uni																		0.25
		MCH clinic, Aizwal													0.96	0	0	0.25		
		MCH, North Bengal																		0
		Medical College & Hospital, Tirunelveli													1.00	1.25	3.00	0.75		
		Medical College Hospital, Bellary														1.67	2.50	0.25		
		Medical College Hospital, Kottayam													0	0	0	0		
		Medical College Hospital, Thrissur													0	0.37	0	0.25		
		Medical college, Amritsar													0	0.50	0	0.50		
		Medical college, Baroda													0	0.76	0.50	0		
		Medical college, Barwani																		0.75
		Medical college, Coimbatore									0.12	0.38	0.25		0.50	0		1.25	0.50	
		Medical college, Faridkot														0.25	0	0		
		Medical college, Guntur													2.75	4.00	3.50	5.25		
		Medical college, Jamnagar													0	0				
		Medical college, Kanpur													0.25	0	0.25	0		
		Medical college, Kurnool															1.25	0.75		
		Medical college, Salem									1.13	0.62	0.75		1.00	0	1.00	1.50		
		Medinipur																0.25	0	
		Mehsana																		0.25
		Mokoko Chung																		1.25
		Murshidabad																0	0.75	
		Muzaffarpur																		0
		Mysore																		1.75
		N.A.R.I., Pune										4.25	2.50	3.49						
Nagpur																		2.25		
Nalgonda																		1.75		
Namakkal																		4.00		
Nasik																		1.75		
Nayagaon																		0		
NB Medical Coll													0.62	0.25	0.50	0.25				

Group	Area	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003		
Pregnant women	Outside major urban areas	Nellore															1.00			
		Nizam Institute of Medical Sciences												28.24	0.50					
		Nizamabad																	1.25	
		Ongole																	3.50	
		Pakyong																	0	
		Palwai																		0.75
		Parbhani																		1.25
		Partapgarh																		0
		Patna																		1.00
		Patna Medical College, Patna													0	0.25	0	0		
		PHC, Daman													0.26	0	0	0.25		
		Pilibhit																		0
		Pithoragarh																		0
		Port Blair																		0
		Primary Health Centre, Pakyong													0	0	0	0		
		Pudukottai																		0.25
		Pune						0.71	3.80											2.88
		Puruliya																		0
		Queen's Mary Hospital, Lucknow													0.25	0	0.75	0		
		Raichur																		2.75
		Raiganj Hospital, Uttar Dinajpur																0.25		
		Raigarh																		1.50
		Rajasthan state													0.08	0.25				
		Rajkot																		0.50
		Rajnandgaon																		0
		Ranchi																		0
		Ranchi Medical College & Hospital,													0	0	0	0		
		Rangaraya MC Hospital, Kakinada															2.00	4.00		
		Rangat													0	0				0
		Ratlam																		0.25
		Ratnagiri																		0.75
		Raxaul																		0
		Rewa																		0
		RIMS & M.C.W., Manipur state									0.81	0.46	0.68	1.32						
		RIMS, Imphal									0.81	0.46	0.68		2.70		0.75	1.75	1.00	
		Rothak district				0	0	0		0.37										
		Rourkela																		0
		S. V. Medical College, Tirupati				0	0.05	0.97	0.79	1.91										
		S.M.S. Medical College, Jaipur													0.29					
		Sadar Hospital, Araria																		0
		Sadar Hospital, Begusarai																		0.25
		Sadar Hospital, Bhagalpur																		0
		Sadar Hospital, Deogarh																		0
		Sadar Hospital, Garhwa																		0
		Sadar Hospital, Jamshehpur															0	0.25		
		Sadar Hospital, Koderma																		0
		Sadar Hospital, Muzaffarpur																		0.25
		Sadar Hospital, Sahibganj																		0
		Sadar Hospital, Sasaran																		0
		Sagar																		0
		Sahibganj																		0
		Salem																		1.50
		Sangareddy																		1.25
		Sangli													5.00	3.00	1.00	4.00	5.00	
		Sasaran																		0.31
		Satara																		3.25
		SCB Medical College, Cuttack													0	0	0.25	0		
		SD Hospital, Durgapur													0.25	0.25	0	0		
		Senapati																		1.25
		Shahdol																		0.25
		Shivpuri																		0.25

Group	Area	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003		
Pregnant women	Outside major urban areas	Siliguri																0		
		Silvassa																	0	
		SMS Medical College, Jaipur district							0											
		Solan																	0	
		Srigangagar																	0	
		Srikakulam																	0.08	
		Srinagar																	0	
		St. John's Medical College & Hospit						0												
		STNM Hospital, Gangtok													0.13	0.25	0	0		
		Surat																		0.50
		Adilabad, Adilabad																		0.75
		Agartala																		0
		Ahmedabad																		0.25
		Aizwal																		1.00
		Akola																		0
		Aligarh																		0
		Along													0.45	0				
		Amritsar																		1.25
		Anakapally																		0.75
		Anantapur																		1.25
		ANC Clinic I, Pune district						1.53												
		ANC Clinic II, Pune district						2.44												
		Andhra Pradesh state													2.20					
		Araria																		0.25
		Asilo Hospital, Mapura									0			0.34	1.73	0.50	1.50	0.50		
		B. J. Medical College, Ahmedabad di					0													
		B. J. Medical College, Pune													2.25	2.75	4.25	2.50		
		Bahraich																		0
		Banda																		0
		Bangalore																		0.75
		Barabanlu																		0.25
		Bardhaman													0.25	0.25	0	0		
		Baroda													0.85					0.50
		Beed																		0.75
		Begusarai																		0.25
		Behrampur																		0.25
		Bellary																		3.00
		Betul																		0
		Bhagalpur																		0.25
		Bhilwara																		0.50
		Bhind																		0
		Bhuj																		0.25
		Bidar																		0.25
		Bijapur																		5.25
		Bilaspur																		2.25
		Bishnupur																		0
		Bomdila																		0
		Bongalgaon, Bongalgaon																		0
		Buldhana																		0
		Burla																		0.25
		Car Nicobar																		0
		Champhai																		3.00
		Chandel																	3.00	
		Chandigarh																		0.25
		Chandrapur																		0.25
		CHC Hospital, Moreh																	3.00	
		CHC, Bishnupur													0.75	2.25				
		CHC, Ponda													0.74	1.00	0.86	0.50		
		CHC, Thoubal													0.75	2.25	0.50	1.00		
		Chennai																		0.50
		Chindwara																		0
		Chittoor																		1.00
		Christian Medical College & Hospita		0.05	0.10	0.04	0.02	0.04	0.08											
		Churachandpur						0.50												8.75
		Civil hospital, Adilabad																	1.25	
		Civil hospital, Anantapur																		2.00
		Civil hospital, Beed																		1.75

Group	Area	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003		
Pregnant women	Outside major urban areas	Civil hospital, Bongalgaon														0				
		Civil hospital, Buldhana															0.25			
		Civil hospital, Car Nicobar												0	0.50	0.75	0			
		Civil hospital, Champai															0.50			
		Civil hospital, Chandrapur															0			
		Civil hospital, Dimapur													2.00		0.75			
		Civil hospital, Diu												0	0	0	0.25			
		Civil hospital, Gadchiroli														0				
		Civil hospital, Hoshiarpur															0.50			
		Civil hospital, Kohima													2.00	0.75	0.75			
		Civil hospital, Ludhiana															0.31			
		Civil hospital, Mokoko Chung													0.71	0.76	0.51	1.75		
		Civil hospital, Parbhani															1.00			
		Civil hospital, Rajkot														1.25	0.50			
		Civil hospital, Silvassa												0	0	0	0.25			
		Civil hospital, Thane														1.50				
		Civil hospital, Tuensang													0.74	4.87	3.51	3.06		
		Civil hospital, Vijayawada															1.25			
		Coimbatore Medical College, Coimbat								0.13	0.25	0.25	1.25							
		Community Health Centre, Garo Hills													0	0.49	0	0.96		
		Community Health Centre, Kakinada													1.99	2.00				
		Community Health Centre, Lungei													0	0	0.75	0.25		
		Community Health Centre, Namakkal													3.25	6.50	3.50	4.25		
		Community Health Centre, Nancowry													0	0				
		Community Health Centre, Nazibad													0	0	0	0		
		Community Health Centre, Rangat															0.25	0.25		
		Cuddalore																	0.25	
		Cuddapah																	1.75	
		Cuttack																	0.25	
		Daman																	0.22	
		Dantewada																	0	
		Deogarh																	0	
		Devangere																	3.25	
		Dewas																	0.25	
		Dimapur																	1.26	
		Dindigul																	0.75	
		District hospital, Bhuj															0.25	0.50		
		District hospital, Bidar																0.51		
		District hospital, Bijapur															4.00	3.25		
		District hospital, Bilaspur																0.50		
		District hospital, Bishnupur															0.50	2.50		
		District hospital, Chandel																	8.00	
		District hospital, Chindwara													0	0	0	0.25		
		District hospital, Chittoor															2.00	1.75		
		District hospital, Churachandpur													3.50	5.25	2.22	3.50		
		District hospital, Dantewada																0		
		District hospital, Devangere																1.25		
District hospital, Dewas																0				
District hospital, Dungarpur															0.25	0.50				
District hospital, Hamirpur													0.24	0.49	0.73	0.25				

Group	Area	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003		
Pregnant women	Outside major urban areas	District hospital, Hassan														1.00				
		District hospital, Himmat Nagar														0.25				
		District hospital, Imphal												0.75	1.88	1.50	1.00			
		District hospital, Jagdalpur												0.94	0.50	0.25	0.50			
		District hospital, Jhalawar															0.25			
		District hospital, Jorhat													0	0	0	0		
		District hospital, Kangra													0.49	0.12		0	0	
		District hospital, Katni																0		
		District hospital, Kinnaur															1.05	0		
		Tamenglong																		1.25
		Tamil Nadu state				0.38	0.32				0.63	0.25	0.41			1.62				0.25
		Tamluk SD Hospital, Midnapur																0.25		
		Tandur																		1.00
		Tezu																		0
		Thane																		2.00
		Thodupuzha																		0.35
		Thoubal																		3.25
		Thrissaur																		0.25
		Tirunelveli																		1.25
		Tripura state										0								
	Tuensang																		8.00	
	Udipi																		1.50	
	Ukhru																		3.25	
	Umaid Hospital & Medical College, J													0	0.25	0.50	0.50			
	Una																		0	
	Uttar Dinajpur																		0.50	
	Uttar Pradesh state													0.20						
	Vani Vials Hospital, Bangalore														0.25	0.50	2.00			
	Vellore																		1.00	
	Victoria Hospital, Bangalore distri								0											
	Vizianagaram																		1.00	
	VS Hospital, Ahmedabad															0.50	0.50			
	Warangal																		0.25	
	Zanana Hospital, Jaipur													0	0	0.25	0			
	Zonal Hospital, Kelong																		0	
	Zonal Hospital, Mandi																		0.50	
	Zonal Hospital, Una																		0.75	
	Major urban areas	Bhagwati Hospital, Mumbai (Bombay)																	2.00	
		Calcutta (1), Calcutta								0	0	0	0.50	0.50						
		Delhi (1), Delhi															0			
		GTB Hospital, New Delhi															0	0		
		J.J. Hospital & Government Medical								2.50		2.38		2.50	3.25	6.50	7.75			
		Kasturba Ghandi, New Delhi																	0.25	
		Kasturba Hospital, New Delhi																0.25		
		KEM Hospital, Mumbai (Bombay)								0		2.25			2.75	2.25	2.75	2.25		
		L. T. Medical College, Mumbai (Bomb)													2.75	2.75	2.00			
Lady Duffrin Hospital, Calcutta																			0	
Madras (1), Madras			0	0																
Madras Medical College, Institute o					0															
Mangolpuri, New Delhi														0.25	0.25	0.25	0.25			
Medical College Hospital, Calcutta														0.75	0					

Group	Area	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003		
Pregnant women	Major urban areas	Micro, K.E.M., Mumbai (Bombay)						0												
		Mumbai (1), Mumbai (Bombay)						0.50	1.71	2.25	2.38	3.00								
		Mumbai (2), Mumbai (Bombay)						1.28					1.20							
		Nair Hospital, Mumbai (Bombay)												3.75	2.00					
		New Delhi				0	0	0.01	0											
		Rajawadi Peripheral Hospital, Mumbai														2.25	3.25			
		RG Kar Medical College, Calcutta																0.50	0.25	
		Safdarjung Hospital, Delhi													0.25	0.25	0.25	0	0.25	
		School of Tropical Medicine, Calcut		0	0	0	0	0	0	0.80										
		SG Hospital, Calcutta															0.50	0	1.25	
Sex workers	Major urban areas	Wadia Hospital, Mumbai (Bombay)						0.65	0.92	0.96	1.67	1.38								
		Calcutta (1), Calcutta									1.20									
		Gaurabai STD clinic, Mumbai (Bombay)												58.80	64.40	58.67	52.26			
		Madras (1), Madras		4.45				7.58												
		Madras (2), Madras																		
		Micro, K.E.M., Mumbai (Bombay)	0.89	8.76	13.20	21.11	33.93		50.99											
		Mumbai (1), Mumbai (Bombay)										51.00					58.67			
		School of Tropical Medicine, Calcut					0	0.87	1.86											
		Sonagachi, Calcutta					0.53	0.72	1.41	1.58	4.18	11.27	11.93	5.30						
		Outside major urban areas	Agra												4.76					
	Ahmedabad state														11.70					
	Autonagar Baina												59.26		20.00					
	Bhiwadi								33.73											
	District hospital, Ujjain		0	0	0	0	0	0	1.94											
	H. J. Doshi Trust Hospital, Rajkot				1.79															
	Kakinada state																42.00			
	Madurai Medical College, Micro. & V		3.11	1.42	4.94	24.23														
	New Civil Hospital, Surat Govt. Med								18.50											
	Pondicherry																3.42			
	Injecting drug users	Major urban areas	Pune	0.49		5.97	21.15	31.58	36.46	46.80	39.92									
Ratlam									1.00	1.00										
Sasson General Hospital, Pune distr									23.71											
Surat									18.52											
Tiruchirapalli																				
Tirupati							25.93													
Trivandrum state																			2.60	
Vasco Da Gama			0				14.00	15.27	28.00	29.00						53.20	50.79			
Vellore region			3.70	5.00	8.50	12.50	19.00	25.00	35.00											
Vijayawada																			51.33	
Injecting drug users	Major urban areas	Visakhapatnam											8.30	13.10	9.70	8.80				
		Calcutta (1), Calcutta							0		1.65	3.52							1.46	
		DDU Hospital, New Delhi														5.00	2.40			
		Madras Drug Addiction Centre, Chenn															26.67	24.56		
		Mumbai (1), Mumbai (Bombay)														23.68				
		New Delhi																	7.20	
SANKALP, Mumbai (Bombay)																41.38				

Group	Area	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003			
Injecting drug users	Major urban areas	School of Tropical Medicine, Calcut				1.92	1.51														
		Outside major urban areas																1.60			
																2.00	2.26				
																		14.80			
																		33.75			
									25.41	47.78	55.71	67.10						50.80			
														0.96	1.55	9.62	2.00				
														13.27	7.56	6.25	7.59	5.83			
											32.00										
														70.73	41.44	44.00	57.20				
						54.54			85.06	85.64	61.05	73.27	76.90						51.60		
														70.33	48.80	64.35	52.00				
			0	0	8.57	39.12	44.75	67.20		55.00	56.00					64.34					
																			0.40		
											3.90					8.00					
														76.10	68.40	75.60	59.60				
															1.20	4.23					
																1.41	1.39				
									1.50	18.90					29.50						
																			24.63		
STI patients	Major urban areas	Army hosp. & base hosp., Delhi										0.08									
		B.Y. L Nair Hospital, Mumbai (Bomba							20.94												
		Balasse Road STD Clinic, Mumbai (Bo				0.22	0.16	1.24	3.24												
		Calcutta (1), Calcutta					0.09	0.13	0.50	1.00	0.86	2.13	5.48								
		Cantonment Board General Hospital,											0								
		Cooper Hospital, Mumbai (Bombay)															7.48				
		Delhi (1), Delhi														3.26	5.83	2.87			
		Gaurabai STD clinic, Mumbai (Bombay)												58.80	64.40						
		GT Hospital, Mumbai (Bombay)																33.69			
		GTB Hospital, New Delhi														3.26	5.83				
		J.J. Hospital & Government Medical								52.02		32.64		34.80	40.00		33.75				
		Khichripur, Delhi												0							
		LNJP Hospital, New Delhi														4.41	2.80	3.60			
		Lok Naik Jayaprakash Narayan Hospit							3.20												
		LTMG Hospital, Mumbai (Bombay)												17.10	9.10						
		Madras Medical College, Institute o			0.27	0.50		0.08	3.05												
		Maulana Azad Hospital, New Delhi								2.40											
		Medical College Hospital, Calcutta													4.00	2.39					
		Mehrauli, Delhi												0							
		Micro, K.E.M., Mumbai (Bombay)				3.22	8.58		23.67												
MT Agarwal Hospital, Mumbai (Bombay)																10.38					
Mumbai (1), Mumbai (Bombay)	1.04	9.68	9.89					13.67	25.55	31.41	26.69	36.58	32.00		33.33		28.50				
Mumbai (2), Mumbai (Bombay)							13.33	29.02									52.30				
Mumbai (3), Mumbai (Bombay)								9.64													
Mumbai Municipal Hospital, Mumbai (19.33						
Safdarjung Hos				0.10	0.09	0.90	0.80						3.60	0.80	2.00	6.40	0.80				

Group	Area	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003
STI patients	Major urban areas	0	0.09	0.03	0										3.24	2.40		
	School of Tropical Medicine, Calcut																	
	Sir J.J. Hospital, Mumbai (Bombay)				13.76	16.36	22.84	24.86										
	Standley Medical College, Madras							3.68	3.78	5.65	5.21		8.00					
	Stanley Medical College, Madras													3.60	5.99	4.00		
	STD Dept., Govt. General Hospital,				4.15	8.85	9.55											
	Outside major urban areas																	
	A. F. Medical College, Pune												2.80	6.80	10.00	10.00		
	Agartala																	2.80
	Agra																	2.00
	Ahmedabad																	8.00
	Ahmadabad								3.98	5.91	6.29							
	Aizwal									1.33								5.20
	Akola																	5.20
	Allahabad																	2.00
	Allappuzha																	0.89
	Almora																	0
	Along																	0
	Ambala																	2.56
	Amritsar																	4.00
	Ara																	0
	Asilo Hospital, Vasco Da Gama								12.64	21.66	16.39							
	Aurangabad								3.75	6.67	14.70							7.60
	Azamgarh																	2.40
	B. J. Medical College, Ahmedabad												6.45	3.20	0.40	7.20	7.60	
	Baina, Vasco Da Gama												22.83	21.89				
	Balasure																	1.20
	Balrampur Hospital, Lucknow												1.20	2.40	2.80	2.00		
	Bangalore																	13.60
	Bareilly																	0.91
	Barmur																	4.80
	Baroda								11.62	9.30	10.22	19.35						4.52
	Basti																	3.20
	Behrampur																	3.20
	Belgaum																	6.00
	Bellary																	18.00
	Bhagalpur																	0
	Bhatinda																	0
	Bhavnagar																	9.68
	Bhiwani																	2.31
	Bhopal									0								1.18
	Bhubaneshwar																	0.80
	Bikaner																	1.60
	Bilaspur																	0.80
	Capital Hospital, Bhubaneshwar												2.86	0		0.80		
	Chamba																	0
	Chandigarh, Chandigarh Union Territ														3.35			
	Chandrapur																	7.60
	Chattarpur																	0
	Chengaipattu																	2.40
	Chennai												8.00	3.60	5.99	4.00		8.80
	Christian Medical College & Hospita	0.40	0.31	2.10	1.20	2.00	3.64	2.25										
	Churachandpur																	13.20
	Civil hospital, Aizwal												1.49	0.76	2.00	4.40		
	Civil hospital, Aurangabad												13.48	20.00	21.60	9.20		
	Civil hospital, Belgaum												40.32	27.34	18.52	23.20		
	Civil hospital, Bellary												21.01	14.06				
	Civil hospital, Bhatinda																	0
	Civil hospital, Car Nicobar								0	0	0							
	Civil hospital, Gulburga												6.00	5.60	12.80	11.60		
	Civil hospital, Hubli												20.80	23.81		16.40		

Group	Area	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003
STI patients	Outside major urban areas													0				
	Civil hospital, Kavarratti															14.80		
	Civil hospital, Khammam																	
	Civil hospital, Kohima												11.11	4.40	6.97	7.42		
	Civil hospital, Mysore												12.75	8.40				
	Civil hospital, Saiha																0	
	Civil hospital, Surat												1.79	13.04	8.00	4.61		
	Coimbatore																	14.72
	Cuttack									0.96								3.20
	Darbanga Medical College Hospital,															1.20		
	Dattenganj Palamu																	0
	Dehradun														6.12			8.00
	Dhanbad																	0.40
	Dhubri																	0.95
	Dibrugrah																	0.95
	District hospital (VIMS), Bellary														8.40	16.40		
	District hospital, Agartala													0.81	1.35	3.20		
	District hospital, Agra												1.60	0.40	0.80	2.00		
	District hospital, Allahabad																	0.80
	District hospital, Almora																	0
	District hospital, Alwar																	0.40
	District hospital, Azamgarh															1.20		
	District hospital, Bareilly														0	0.96		
	District hospital, Barmur														2.88	1.20		
	District hospital, Bhavnagar														7.39	4.74		
	District hospital, Bilaspur												0	0	0	0.80		
	District hospital, Chamba														1.18	0		
	District hospital, Chattarpur														0	0		
	District hospital, Churachandpur												2.00	13.20	11.76	11.20		
	District hospital, Dehradun														8.16	0.80		
	District hospital, Durg																	1.76
	District hospital, Ghaziabad														3.17	2.40		
	District hospital, Godhara														0.71	0.88		
	District hospital, Gonda												3.26	0.80	0.40	0.86		
	District hospital, Jabalpur														1.43	0.65		
	District hospital, Jhansi																	0.79
	District hospital, Junagarh																	0.50
	District hospital, Katihar														7.56	6.67		
	District hospital, Korba														1.60	0		
	District hospital, Kullu														0	0		
	District hospital, Malda																	0
	District hospital, Muzaffar																	0.40
	District hospital, Nahan												0.40	0.80	0.79	0		
	District hospital, Palampur														2.08	2.40		
	District hospital, Pasighat												0	0	0	0		
	District hospital, Seoni														2.80	1.22		
	District hospital, Tehri Garhwal																	2.33
	District hospital, Uttarkashi																	0
	District hospital, Varanasi												1.60	2.00	0.42	0.40		
	Dr. S. N. Medical College, Jodhpur									7.24	5.67	5.44	11.16					
	Duncan Hospital, Raxaul												1.61	3.56	2.40	6.37		
	Durg																	1.60

Group	Area	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003		
STI patients	Outside major urban areas	East Delhi																7.30		
		Emakulam																	4.40	
		Farrukhabad																	1.88	
		Gandhinagar Hospital, Jammu												1.84	1.16	0.80	1.20			
		Gangtok							0	0									0	
		Gaya																		2.00
		Gaya Medical College, Gaya													0.40	0	0.40	0.40		
		GB Pant Hospital, Port Blair								0	1.33	0	0		1.27	0.40	1.20	1.20		
		General hospital, Ambala																0	1.05	
		General hospital, Chandigarh Union								0	0	0.49	2.55							
		General hospital, Karaikal															2.80	1.20		
		General Hospital, Sector 16, Chand								0	0	0.49	2.55		4.08	1.98	3.36	0.82		
		General hospital, Sirsa															3.77	1.20		
		GH, Chandigarh																		0.80
		Ghaziabad																		0
		GMC, Srinagar															0	0.40		
		Godhara																		6.56
		Gonda																		0.80
		Gorakhpur																		2.18
		Government hospital, Akola																	10.40	
		Government hospital, Amritsar													0	1.60	1.20	3.60		
		Government hospital, Balasore																	0.40	
		Government hospital, Chandrapur													4.80	7.20	5.60	4.40		
		Government hospital, Gurgaon															2.86	0.80		
		Government hospital, Gwalior													1.22	0	0.68	0.66		
		Government hospital, Jalgaon																	11.60	
		Government hospital, Koraput																	1.20	
		Government hospital, Latur													16.00	15.20	18.40	5.60		
		Government hospital, Nagpur													16.00	20.00	25.11	17.27		
		Government hospital, Naharlagun													0	0	2.04	0.79		
		Government hospital, Ramanathapuram																	1.27	
		Government hospital, Sambalpur																	0.56	
		Government hospital, Tanjore																	9.60	
		Government hospital, Thane																	9.20	
		Government HQS Hospital, Tiruchirap													16.30	34.80	48.00	22.03		
		Government Medical College, Sangli													50.25	44.80	41.82	22.80		
Government Rajaji Hospital, Madurai		0.10	0.34	0.94	2.42	2.76	4.15	5.22												
GS Hospital, Jamnagar															0	3.66	0.61			
Gulburga																		10.40		
Gurgaon																		0.40		
Guwahati																		0.58		
Gwalior																		1.61		
Hamirpur																		0.40		
Headquarter Hospital, Puri														0.40	1.63	0.40				
Hoogly																		0.47		
Hoshangabad																		0.62		
Hospico Hospital, Margao													16.00	5.05	13.04	15.00				
Hubli																		13.60		

Group	Area	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003
STI patients	Outside major urban areas														23.20	13.20		
	KR Hospital, Mysore																	10.40
	Kurnool																	1.20
	Laheriasarai																	6.40
	Latur																	0
	Lucknow																	24.80
	Madurai					4.43			6.52	8.64	10.00	22.40						
	Madurai Medical College, Madurai												23.20	10.40	16.80	15.60		
	Madurai Medical College, Micro. & V	0.29	1.04	1.34	2.49	6.63	6.11	4.85										
	Mangalore																	11.40
	Medical College Hospital, Behrampur													4.80	2.63	4.80		
	Medical college, Allappuzha														0.72	3.33		
	Medical college, Bankura												0.80	0.45	0.40	1.25		
	Medical college, Bhopal													0.40	1.71	1.78		
	Medical college, Burdwan												0	0.44	0.40	0.40		
	Medical College, Dibrugrah												0.40		0.83	1.20		
	Medical College, Guwahati												3.80	2.44	0.48	1.79		
	Medical College, Indore												4.35	2.40	2.40	6.87		
	Medical College, Kozhikode												3.20	4.35	7.73	7.83		
	Medical College, Panaji														11.01	13.46		
	Medical College, Patiala													2.44	0.40	1.23		
	Medical College, Rohtak												2.56	5.26	2.66	1.12		
	Medical College, Trivandrum												3.88	1.97	5.22	5.05		
	Medical College, Udaipur												4.40	3.20	2.80	8.00		
	Medical College, Visakhapatnam								7.80				21.60	29.51	30.00	38.40		
	Microbiology department, Manipal								1.67									
	Mirzapur																	4.86
	MJM Hospital, Coochbihar													2.59	1.96	0		
	Moradabad																	0.40
	Morena																	0.82
	MR Bagur Hospital, Hoogly														2.78	0.80		
	Muzaffar Nagar																	0
	Muzaffarpur																	0.80
	Mysore																	21.60
	Nagpur								6.75	3.25	5.75	11.75						21.20
	Nahan																	0.80
	Naharlagun																	0.64
	Ongole																	12.80
	Osmania Medical College, Hyderabad					1.33	2.60	2.64	4.69	4.44			34.77	27.60	32.00	41.60		
	Palampur																	1.60
	Pasighat																	0
	Patiala																	0.81
	Patilputra Medical College & Hospi														0.61	0.38		
	Patna																	2.80
	Patna Medical College, Patna													0.80	0	0		
	PBM Hospital, Bikaner														2.55	0.40		
	PGI, Chandigarh												0		0	6.74	2.82	
	PGI, Chandigarh Union Territory												0	1.56				
	Port Blair																	0
	Prothrapur																	5.20
	Pt. JNM Medical College, Raipur												3.96	0	1.60	2.83		
	Pune	0.21	1.42	3.02	4.80	8.84			18.40	20.00	18.50	19.20	21.70	20.30	21.40			5.20
	Puri																	0

Group	Area	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003		
STI patients	Outside major urban areas	Raipur																0.80		
		Rajahmundry																	30.40	
		Ramanathapuram																	4.35	
		Ranchi																	0	
		Ranchi Medical College & Hospital,												1.35	0.40	0	0			
		Raxaul																	7.60	
		Rohtak																	1.48	
		Roing																	0	
		S. V. Medical College, Tirupati	0	0.82	1.17	1.86	1.67	13.75	6.19	5.75	6.25				9.60	30.00	23.60	12.97		
		S.M.S. Medical College, Jaipur													6.00	3.20	12.40	6.80		
		Sadar Hospital, Dattenganj Palamu																	0.38	
		Saiha																		0
		Sambalpur																		0.40
		Sangareddy																		3.20
		Sangli																		20.00
		Satna																		3.85
		SCB Medical College, Cuttack														2.00	3.81	1.20		
		Selam																		14.80
		Seoni																		1.20
		Silchar																		0
		Simla									0	0.34	0							0
		Sirsa																		0.80
		Srinagar																		0.80
		SSG Hospital, Baroda																	5.26	
		St. John's Medical College & Hospit													8.40	16.78				
		State AIDS cell, Pondicherry									7.60	8.00	6.00		7.20	2.00	4.17			
		STNM Hospital, Gangtok														0	0	0.96		
		Sultanpur																		0.40
		Surat									13.88	19.39	19.31	20.62						5.76
		Surendranagar																		1.56
		Tanjore																		11.20
		Tehri Garhwal																		0
		Tezpur																		0
		Thane																		8.00
		Tiruchirapalli																		15.20
		Tirunelveli																		22.00
		Tirupati																		39.20
		Trivandrum																		4.00
		Tura																	0	
		Udaipur																		7.60
		Ujjain																		2.35
		Uttarkashi																		0
		Varanasi																		4.40
		Vellore																		10.40
		Victoria Hospital, Bangalore									7.34	7.31					0	14.80		
		Vijayawada																		33.20
		Visakhapatnam																		35.60
		Hyderabad																		31.60
		I. G. Medical College, Simla													0.39	0.39	0	0.52		
		Imphal									4.81	3.91	8.17	4.85						6.00
Indore								2.56										7.29		
J. N. Hospital, Imphal													6.28	10.80	16.79	8.80				
Jabalpur																		0		
Jaipur									1.36		1.13							12.40		
Jalgaon																		4.00		
Jammu																		0.80		
Jantia Hills Community Health Centr															0	0				
Jhansi																		0		
JIPMER, Pondicherry														9.60	17.53	11.63				
JLN Medical College & Hospital, Aim																		8.40		
Karnataka Medical College, Hubli									20.50	16.91										
Karur																		18.80		

Group	Area	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	
STI patients	Outside major urban areas	Kasturba Medical College, Mangalore												15.54	10.80	24.00			
		Katihar																7.60	
		Khammam																12.80	
		KIMS, Hubli								20.50	16.91					17.60			
		Kohima									3.00							2.43	
		Korba																0	
		Kota																0	
Men having sex with men	Major urban areas	HAMSAFAR, Mumbai (Bombay)													23.94	23.60			
		Mumbai (1), Mumbai (Bombay)						10.32							14.00	17.21	23.60		
		Mumbai (2), Mumbai (Bombay)						15.90											
	Outside major urban areas	Chennai																1.20	
		Goa		0.37															
		ICWO, Tamil Nadu state																3.60	
		Madhya Pradesh state		0															
Tuberculosis patients	Major urban areas	Calcutta (1), Calcutta					0	0	1.00	0.50	3.20	3.40							
		J.J. Hospital & Government Medical			3.79	3.96	7.07	8.41											
		Madras (1), Madras										1.50	2.86						
		Sir J.J. Hospital, Mumbai (Bombay)		2.40	2.30	3.90	7.90	10.60	10.60	11.00	10.70	16.80	23.00	30.00	33.00				
		TB Research Centre, Madras	0		0		0.35	0.76											
	Outside major urban areas	Dr. S. N. Medical College, Jodhpur										15.71	21.43	11.16					
		DTC, Vellore						4.46											
		GBTs, Pondicherry Union Territory						0.71											
		Government hospital, TB clinic, Pe												6.00	2.80				
		Government hospital, TB clinic, Ta												17.60	16.40				
		GTBS, Vellore						2.00											
Tuberculosis patients	Outside major urban areas	Kerala state						3.20											
		Manipur state							11.50	4.00	6.17	11.86							
		Perundurai									3.25								
		Pondicherry Union Territory						0.90											
		Pune						1.10		14.30	12.10	18.50	21.50						
		Regional Medical Research Centre, B							0										
		Sassoon General Hospital, Pune				3.20			5.06	10.29									
		Tambaram Hospital, Tambaram						0.98											
		Tamil Nadu state									2.25	3.60	4.39	7.00	9.60				
		TB & Chest Dept., Gov. Medical Coll									5.19				20.06				
		TB center, Imphal								3.20				14.80					
Vellore						1.32													
Yavatmal/ rural																4.50			