

2004  
Update



## United Republic of Tanzania

**EPIDEMIOLOGICAL FACT SHEETS**  
ON HIV/AIDS AND SEXUALLY TRANSMITTED INFECTIONS



Joint United Nations Programme on HIV/AIDS

**UNAIDS**

UNHCR • UNICEF • WFP • UNDP • UNFPA  
UNODC • ILO • UNESCO • WHO • WORLD BANK



World Health  
Organization

## HIV/AIDS estimates

In 2003 and during the first quarter of 2004, UNAIDS and WHO worked closely with national governments and research institutions to recalculate current estimates on people living with HIV/AIDS. These calculations are based on the previously published estimates for 1999 and 2001 and recent trends in HIV/AIDS surveillance in various populations. A methodology developed in collaboration with an international group of experts was used to calculate the new estimates on prevalence and incidence of HIV and AIDS deaths, as well as the number of children infected through mother-to-child transmission of HIV. Different approaches were used to estimate HIV prevalence in countries with low-level, concentrated or generalised epidemics. The current estimates do not claim to be an exact count of infections. Rather, they use a methodology that has thus far proved accurate in producing estimates that give a good indication of the magnitude of the epidemic in individual countries. However, these estimates are constantly being revised as countries improve their surveillance systems and collect more information.

Adults in this report are defined as women and men aged 15 to 49. This age range covers people in their most sexually active years. While the risk of HIV infection obviously continues beyond the age of 50, the vast majority of those who engage in substantial risk behaviours are likely to be infected by this age. The 15 to 49 range was used as the denominator in calculating adult HIV prevalence.

### Estimated number of adults and children living with HIV/AIDS, end of 2003

These estimates include all people with HIV infection, whether or not they have developed symptoms of AIDS, alive at the end of 2003:

Adults and children	1,600,000		
Low estimate	1,200,000		
High estimate	2,300,000		
Adults (15-49)	1,500,000	Adult rate (%)	8.8
Low estimate	1,100,0	Low estimate	6.4
High estimate	2,000,000	High estimate	11.9
Children (0-15)	140,000		
Low estimate	85,000		
High estimate	230,000		
Women (15-49)	840,000		
Low estimate	610,000		
High estimate	1,100,000		

### Estimated number of deaths due to AIDS

Estimated number of adults and children who died of AIDS during 2003:

Deaths in 2003	160,000
Low estimate	110,000
High estimate	230,000

### Estimated number of orphans

Estimated number of children who have lost their mother or father or both parents to AIDS and who were alive and under age 17 at the end of 2003:

Current living orphans	980,000
Low estimate	660,000
High estimate	1,400,000

## Assessment of the epidemiological situation 2004

HIV information among antenatal clinic attendees is available from Tanzania since the mid-1980s. In Dar es Salaam, the major urban area, HIV prevalence among antenatal clinic attendees tested increased from 4 percent in 1986 to 15 percent in 1999. In 2002, HIV prevalence among attending ANC clinics in Dar es Salaam was 11.5 percent. In 2002, the median HIV prevalence at 24 antenatal care clinics in 6 regions was 8.1 percent. The highest prevalence was reported in Mbeya region (median 17.2 percent) followed by Dar es Salaam.

Information on HIV prevalence among sex workers in Dar es Salaam is available since the mid-1980s. HIV prevalence among sex workers tested increased from 29 percent in 1986 to 49.5 percent in 1993. In 2001, nearly 70 percent of sex workers tested in Mbeya were HIV positive.

In Dar es Salaam, HIV prevalence among STI clinic patients tested increased from 13 percent in 1986 to 24 percent in 1994. In 2000 and 2001, over 40 percent of STI patients tested positive for HIV in Dar es Salaam. Outside of Dar es Salaam, the majority of data on HIV prevalence among STI clinic patients comes from Mbeya. HIV prevalence among STI patients tested in Mbeya increased from 22.5 percent in 1988 to 27 percent in 1997.

## UNAIDS/WHO Working Group on Global HIV/AIDS and STI Surveillance

Global Surveillance of HIV/AIDS and sexually transmitted infections (STIs) is a joint effort of WHO and UNAIDS. The UNAIDS/WHO Working Group on Global HIV/AIDS and STI Surveillance, initiated in November 1996, guides respective activities. The primary objective of the Working Group is to strengthen national, regional and global structures and networks for improved monitoring and surveillance of HIV/AIDS and STIs. For this purpose, the Working Group collaborates closely with national AIDS programmes and a number of national and international experts and institutions. The goal of this collaboration is to compile the best information available and to improve the quality of data needed for informed decision-making and planning at national, regional, and global levels. The Epidemiological Fact Sheets are one of the products of this close and fruitful collaboration across the globe.

Within this framework, the Fact Sheets collate the most recent country-specific data on HIV/AIDS prevalence and incidence, together with information on behaviours (e.g. casual sex and condom use) which can spur or stem the transmission of HIV.

Not unexpectedly, information on all of the agreed upon indicators was not available for many countries in 2003. However, these updated Fact Sheets do contain a wealth of information which allows identification of strengths in currently existing programmes and comparisons between countries and regions. The Fact Sheets may also be instrumental in identifying potential partners when planning and implementing improved surveillance systems.

The fact sheets can be only as good as information made available to the UNAIDS/WHO Working Group on Global HIV/AIDS and STI Surveillance. Therefore, the Working Group would like to encourage all programme managers as well as national and international experts to communicate additional information to them whenever such information becomes available. The Working Group also welcomes any suggestions for additional indicators or information proven to be useful in national or international decision-making and planning.

## Basic indicators

For consistency reasons the data used in the table below are taken from official UN publications.

DEMOGRAPHIC DATA	YEAR	ESTIMATE	SOURCE
Total population (thousands)	2004	37,671	UN population division database
Female population aged 15-24 (thousands)	2004	4,042	UN population division database
Population aged 15-49 (thousands)	2004	17,629	UN population division database
Annual population growth rate (%)	1992-2002	2.6	UN population division database
% of population in urban areas	2003	34.9	UN population division database
Average annual growth rate of urban population	2000-2005	4.9	UN population division database
Crude birth rate (births per 1,000 pop.)	2004	38.4	UN population division database
Crude death rate (deaths per 1,000 pop.)	2004	18.2	UN population division database
Maternal mortality rate (per 100,000 live births)	2000	1500	WHO (WHR2004)/UNICEF
Life expectancy at birth (years)	2002	46.5	World Health Report 2004, WHO
Total fertility rate	2002	5.2	World Health Report 2004, WHO
Infant mortality rate (per 1,000 live births)	2000	109	World Health Report 2004, WHO
Under 5 mortality rate (per 1,000 live births)	2000	156	World Health Report 2004, WHO

SOCIO-ECONOMIC DATA	YEAR	ESTIMATE	SOURCE
Gross national income, ppp, per capita (Int.\$)	2002	550	World Bank
Gross domestic product, per capita % growth	2001-2002	3.6	World Bank
Per capita total expenditure on health (Int.\$)	2001	26	World Health Report 2004, WHO
General government expenditure on health as % of total expenditure on health	2001	46.7	World Health Report 2004, WHO
Total adult illiteracy rate	2000	25	UNESCO
Adult male illiteracy rate	2000	16.1	UNESCO
Adult female illiteracy rate	2000	33.5	UNESCO
Gross primary school enrolment ratio, male	2000/2001	63	UNESCO
Gross primary school enrolment ratio, female	2000/2001	63	UNESCO
Gross secondary school enrolment ratio, male	2000/2001	6	UNESCO
Gross secondary school enrolment ratio, female	2000/2001	5	UNESCO

### Contact address

UNAIDS/WHO Working Group on Global  
HIV/AIDS and STI Surveillance  
20, Avenue Appia  
CH - 1211 Geneva 27  
Switzerland  
Fax: +41-22-791-4834

email: [hivstrategicinfo@who.int](mailto:hivstrategicinfo@who.int) or  
[estimates@unaids.org](mailto:estimates@unaids.org)

website: <http://www.who.int/hiv>  
<http://www.unaids.org>

Extracts of the information contained in these fact sheets may be reviewed, reproduced or translated for research or private study but not for sale or for use in conjunction with commercial purposes. Any use of information in these fact sheets should be accompanied by the following acknowledgment "UNAIDS/WHO epidemiological fact sheets on HIV/AIDS and Sexually Transmitted Infections, 2004 Update".

## HIV prevalence in different populations

This section contains information about HIV prevalence in different populations. The data reported in the tables below are mainly based on the HIV database maintained by the United States Bureau of the Census where data from different sources, including national reports, scientific publications and international conferences are compiled. To provide a simple overview of the current situation and trends over time, summary data are given by population group, geographical area (Major Urban Areas versus Outside Major Urban Areas), and year of survey. Studies conducted in the same year are aggregated and the median prevalence rates (in percentages) are given for each of the categories. The maximum and minimum prevalence rates observed, as well as the total number of surveys/sentinel sites, are provided with the median, to give an overview of the diversity of HIV-prevalence results in a given population within the country. Data by sentinel site or specific study from which the medians were calculated are printed at the end of this fact sheet.

The differentiation between the two geographical areas Major Urban Areas and Outside Major Urban Areas is not based on strict criteria, such as the number of inhabitants. For most countries, Major Urban Areas were considered to be the capital city and - where applicable - other metropolitan areas with similar socio-economic patterns. The term Outside Major Urban Areas considers that most sentinel sites are not located in strictly rural areas, even if they are located in somewhat rural districts.

### HIV sentinel surveillance\*


Group	Area		1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003			
Pregnant women	Major urban areas	N-Sites		1.00	1.00	1.00	2.00	1.00	4.00	1.00	3.00	1.00	1.00		3.00	2.00		4.00				
		Minimum		7.80	8.93	8.95	9.12	11.03	10.61	13.81	7.30	12.99	13.73		14.10	10.10		10.00				
		Median		7.80	8.93	8.95	9.72	11.03	15.24	13.81	12.17	12.99	13.73		15.30	12.20		11.46				
		Maximum		7.80	8.93	8.95	10.33	11.03	16.10	13.81	13.70	12.99	13.73		18.10	14.30		16.29				
	Outside major urban areas	N-Sites	2.00	12.00	11.00	10.00	16.00	23.00	24.00	19.00	22.00	17.00	18.00	11.00	19.00	28.00	1.00	21.00				
		Minimum	0.26	0.53	0.44	0.29	0.68	0.50	0.77	0.30	0	0	0	10.20	0.50	0	15.00	0.49				
		Median	3.14	4.84	9.09	8.65	7.80	8.00	10.76	15.00	9.60	9.10	14.25	14.50	12.10	12.75	15.00	6.08				
		Maximum	6.02	11.00	21.21	22.24	21.03	30.36	27.23	27.50	33.30	25.90	36.00	24.90	40.10	32.10	15.00	17.96				
	Sex workers	Major urban areas	N-Sites		2.00	2.00	1.00	1.00		1.00												
			Minimum		35.71	26.71	44.97	42.86		49.46												
			Median		38.90	35.22	44.97	42.86		49.46												
			Maximum		42.08	43.73	44.97	42.86		49.46												
Outside major urban areas		N-Sites	1.00	5.00				1.00	1.00								1.00	2.00				
		Minimum	14.80	7.50				39.81	60.87								44.60	39.00				
		Median	14.80	19.50				39.81	60.87								44.60	53.50				
		Maximum	14.80	31.90				39.81	60.87								44.60	68.00				
Injecting drug users		STI patients	Major urban areas	N-Sites	2.00	2.00	2.00	2.00	6.00		1.00			2.00			1.00	1.00				
				Minimum	13.37	21.83	28.32	21.05	14.35		23.97				5.08			46.00	41.50			
				Median	20.72	24.81	33.99	22.77	18.88		23.97				14.75			46.00	41.50			
				Maximum	28.07	27.78	39.66	24.49	40.38		23.97				24.42			46.00	41.50			
	Outside major urban areas		N-Sites	2.00	1.00	2.00	1.00	1.00	2.00					9.00								
			Minimum	20.00	22.50	21.35	27.50	32.00	5.16					17.00								
			Median	33.34	22.50	22.48	27.50	32.00	19.73					27.00								
			Maximum	46.67	22.50	23.60	27.50	32.00	34.30					43.00								
	Men having sex with men		Tuberculosis patients	Major urban areas	N-Sites		1.00			1.00	1.00		1.00	3.00								
					Minimum		35.83			65.62	55.22		54.79	52.37								
					Median		35.83			65.62	55.22		54.79	52.37								
					Maximum		35.83			65.62	55.22		54.79	53.14								
Outside major urban areas		N-Sites		1.00			1.00	8.00	5.00	9.00				17.00				1.00				
		Minimum		6.34			54.00	19.73	9.62	16.48				21.55				32.00				
		Median		6.34			54.00	31.37	21.67	26.30				34.14				32.00				
		Maximum		6.34			54.00	54.05	33.00	59.04				70.54				32.00				

\*Detailed data by site can be found in the Annex.


## Maps & charts

Mapping the geographical distribution of HIV prevalence among different population groups may assist in interpreting both the national coverage of the HIV surveillance system as well in explaining differences in levels of prevalence. The UNAIDS/WHO Working Group on Global HIV/AIDS and STI Surveillance, in collaboration with the WHO Public Health Mapping Team, Communicable Diseases, is producing maps showing the location and HIV prevalence in relation to population density, major urban areas and communication routes. For generalized epidemics, these maps show the location of prevalence of antenatal surveillance sites.

Trends in antenatal sentinel surveillance for higher prevalence countries, or in prevalence among selected populations for countries with concentrated epidemics, are a new addition. These are presented for those countries where sufficient data exist.



### Trends in HIV prevalence among antenatal clinic attendees



Median prevalence and ranges are shown in areas with more than one sentinel site.

The boundaries and names shown and the designations used on the map do not imply the expression of any opinion whatsoever on the part of the World Health Organization concerning the legal status of any country, territory, city or area or of its authorities, or concerning the delimitation of its frontiers or boundaries. Dotted lines on maps represent approximate border lines for which there may not yet be full agreement. WHO 2004, all rights reserved.

## Reported AIDS cases

Following WHO and UNAIDS recommendations, AIDS case reporting is carried out in most countries. Data from individual AIDS cases are aggregated at the national level and sent to WHO. However, case reports come from surveillance systems of varying quality. Reporting rates vary substantially from country to country and low reporting rates are common in developing countries due to weaknesses in the health care and epidemiological systems. In addition, countries use different AIDS case definitions. A main disadvantage of AIDS case reporting is that it only provides information on transmission patterns and levels of infection approximately 5-10 years in the past, limiting its usefulness for monitoring recent HIV infections.

Despite these caveats, AIDS case reporting remains an important advocacy tool and is useful in estimating the burden of HIV-related morbidity as well as for short-term planning of health care services. AIDS case reports also provide information on the demographic and geographic characteristics of the affected population and on the relative importance of the various exposure risks. In some situations, AIDS reports can be used to estimate earlier HIV infection patterns using back-calculation. AIDS case reports and AIDS deaths have been dramatically reduced in industrialized countries with the introduction of Anti-Retroviral Therapy (ART).

1979	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998
				3	109	292	1121	2937	4839	5096	11106	18692	15871	13506	6096	4722	8426	10592	8675
1999	2000	2001	2002	2003	Total	UNK	Date of last report												
8850	11673				132606		12/11/2001												

## Curable sexually transmitted infections (STIs)

The predominant mode of transmission of both HIV and other STIs is sexual intercourse. Measures for preventing sexual transmission of HIV and STIs are the same, as are the target audiences for interventions. In addition, strong evidence supports several biological mechanisms through which STIs facilitate HIV transmission by increasing both HIV infectiousness and HIV susceptibility. Thus, detection and treatment of individuals with STIs is an important part of an HIV control strategy. In summary, if the incidence/prevalence of STIs is high in a country, then there is the possibility of high rates of sexual transmission of HIV. Monitoring trends in STIs provides valuable insight into the likelihood of the importance of sexual transmission of HIV within a country, and is part of second generation surveillance. These trends also assist in assessing the impact of behavioural interventions, such as delaying sexual debut, reducing the number of sex partners and promoting condom use.

Clinical services offering STI care are an important access point for people at high risk for both STIs and HIV. Identifying people with STIs allows for not only the benefit of treating the STI, but for prevention education, HIV testing, identifying HIV-infected persons in need of care, and partner notification for STIs or HIV infection. Consequently, monitoring different components of STI prevention and control can also provide information on HIV prevention and control activities within a country.

### STI syndromes

Reported cases	1996	1997	1998	1999	2000	2001	2002	2003	Incidence 2003
Urethral discharge						30019			
Genital Ulcer						34755			

Comments:

Source:

### Syphilis prevalence, women

Percent of blood samples taken from pregnant women aged 15-49 that test positive for syphilis - positive reaginic and treponemal test - during routine screening at selected antenatal clinics.

Year	Area	Rate	Range

Comments:

Source:

### Estimated prevalence of curable STIs among female sex workers

#### - Chlamydia

Year	Area	Rate	Range

Comments:

Source:

#### - Gonorrhoea

Year	Area	Rate	Range

Comments:

Source:

Estimated prevalence of curable STIs among female sex workers (continued)**- Syphilis**

Year	Area	Rate	Range
------	------	------	-------

Comments:

Source:

**- Trichomoniasis**

Year	Area	Rate	Range
------	------	------	-------

Comments:

Source:

## Health service and care indicators

HIV prevention strategies depend on the twin efforts of care and support for those living with HIV or AIDS, and targeted prevention for all people at risk or vulnerable to the infection. It is difficult to capture such a large range of activities with one or just a few indicators. However, a set of well-established health care indicators may help to identify general strengths and weaknesses of health systems. Specific indicators, such as access to testing and blood screening for HIV, help to measure the capacity of health services to respond to HIV/AIDS - related issues.

### Access to health care

Indicators	Year	Estimate	Source
% of population with access to health services - total			
% of population with access to health services - urban			
% of population with access to health services - rural			
Contraceptive prevalence rate (%)	1999	24.2	UNICEF/UNPOP
Percentage of contraceptive users using condoms			
% of births attended by skilled health personnel	2000	35.8	WHO
% of 1-yr-old children fully immunized - DPT	2002	89	WHO/UNICEF
% of 1-yr-old children fully immunized - Measles	2001	83	WHO/UNICEF
% of ANC clinics where HIV testing is available			

### Number of adults (15-49) with advanced HIV infection receiving ARV therapy as of June 2004

#### Adults on treatment

Number: 1,650

Source: WHO

### Estimated number of adults (15-49) in need of treatment in 2003

#### Adults needing treatment

Number: 260,000

Source: WHO/UNAIDS

### Coverage of HIV testing and counselling

Number of public and NGO services providing testing and counselling services.

Year	Area	N=
------	------	----

Comments:

Source:

## Knowledge and behaviour

In most countries the HIV epidemic is driven by behaviours (e.g.: multiple sexual partners, injecting drug use) that expose individuals to the risk of infection. Information on knowledge and on the level and intensity of risk behaviour related to HIV/AIDS is essential in identifying populations most at risk for HIV infection and in better understanding the dynamics of the epidemic. It is also critical information in assessing changes over time as a result of prevention efforts. One of the main goals of the 2nd generation HIV surveillance systems is the promotion of a standard set of indicators defined in the National Guide (Source: National AIDS Programmes, A Guide to Monitoring and Evaluation, UNAIDS/00.17) and regular behavioural surveys in order to monitor trends in behaviours and to target interventions.

The indicators on knowledge and misconceptions are an important prerequisite for prevention programmes to focus on increasing people's knowledge about sexual transmission, and, to overcome the misconceptions that act as a disincentive to behaviour change. Indicators on sexual behaviour and the promotion of safer sexual behaviour are at the core of AIDS programmes, particularly with young people who are not yet sexually active or are embarking on their sexual lives, and who are more amenable to behavioural change than adults. Finally, higher risk male-male sex reports on unprotected anal intercourse, the highest risk behaviour for HIV among men who have sex with men.

### Knowledge of HIV prevention methods

Prevention indicator: Percentage of young people 15-24 who both correctly identify two ways of preventing the sexual transmission of HIV and who reject three misconceptions about HIV transmission.

Year	Male	Female
1999	29	26

Comments:

Source: DHS

### Reported condom use at last higher risk sex (young people 15-24)

Prevention indicator: Proportion of young people reporting the use of a condom during sex with a non-regular partner.

Year	Male	Female
1999	31	21

Comments:

Source: DHS

### Age-mixing in sexual partnerships among young women

The proportion of young women who have had sex in the last 12 months with a partner who is 10 or more years older than themselves.

Year	Area	Age group	Male	Female	All
------	------	-----------	------	--------	-----

Comments:

Source:

### Reported non-regular sexual partnerships

Prevention indicator: Proportion of young people 15-24 having at least one sex partner other than a regular partner in the last 12 months.

Year	Male	Female
1999	87	40

Comments:

Source: DHS

Knowledge and behaviour (continued)Ever used a condom

Percentage of people who ever used a condom.

Year	Area	Age group	Male	Female	All
------	------	-----------	------	--------	-----

Comments:

Source:

Adolescent pregnancy

Percentage of teenagers 15-19 who are mothers or pregnant with their first child.

Year	Percentage
------	------------

Comments:

Source:

Age at first sexual experience

Proportion of 15-19 year olds who have had sex before age 15.

Year	Male	Female
1999	24	15

Comments:

Source: DHS

## Prevention indicators

Male and female condoms are the only technology available that can prevent sexual transmission of HIV and other STIs. Persons exposing themselves to the risk of sexual transmission of HIV should have consistent access to high quality condoms. AIDS Programs implement activities to increase both availability of and access to condoms. These activities should be monitored and have resources directed to problem areas. The indicator below highlights the availability of condoms. However, even if condoms are widely available, this does not mean that individuals can or do access them.

### Condom availability nationwide

Total number of condoms available for distribution nationwide during the preceding 12 months, divided by the total population aged 15-49.

Year	N	Rate
------	---	------

**Comments:**

**Source:**

### Prevention of mother-to-child transmission (MTCT) nationwide

Percentage of women who were counselled during antenatal care for their most recent pregnancy, accepted an offer of testing and received their test results, of all women who were pregnant at any time in the preceding two years.

Year	N	Rate
2000	22489	31.1

**Comments:** This data is collected only from five public health institutions where pilot projects are ongoing.

**Source:** Ministry of Health.

Blood safety programs aim to ensure that the majority of blood units are screened for HIV and other infectious agents. This indicator gives an idea of the overall percentage of blood units that have been screened to high enough standards that they can confidently be declared free of HIV.

### Screening of blood transfusions nationwide

Percentage of blood units transfused in the last 12 months that have been adequately screened for HIV according to national or WHO guidelines.

Year	N	Rate
2000	128,595	100

**Comments:**

**Source:** Ministry of Health.

## Sources

Data presented in this Epidemiological Fact Sheet come from several sources, including global, regional and country reports, published documents and articles, posters and presentations at international conferences, and estimates produced by UNAIDS, WHO and other United Nations agencies. This section contains a list of the more relevant sources used for the preparation of the Fact Sheet. Where available, it also lists selected national Web sites where additional information on HIV/AIDS and STI are presented and regularly updated. However, UNAIDS and WHO do not warrant that the information in these sites is complete and correct and shall not be liable whatsoever for any damages incurred as a result of their use.

Ali, A. K., O. J. Khatib, W. Osei, et al. 1992 Sentinel Surveillance for HIV Infection: Five Years Period VII International Conference on AIDS in Africa, Yaounde, Cameroon, 12/8-12, Poster T.P.036.

Ahmed, H. J., J. Mbwana, C. Guerino, et al. 2001 Etiology of Genital Ulcer Diseases and Prevalence of HIV Co-Infection in a Sexually Transmitted Diseases Clinic in Dar es Salaam. Presented at International Congress of Sexually Transmitted Infections, ISSTD/ISTD, Berlin, Germany, 24-27 June 2001, Abstract in International Journal of STD & AIDS, vol. 12, suppl. 2, p. 133.

Chum, H. J., R. J. O'Brien, T. M. Chonde, et al. 1996 An Epidemiological Study of Tuberculosis and HIV Infection in Tanzania, 1991-1993 AIDS, vol. 10, no. 3, pp. 299-309.

Clift, S., Z. Kanga, L. Ndeki, et al. 2001 Baseline Prevalence of HIV Infection and Other STIs and Their Associated Risk Factors in 2 Gold Mining Communities of the ... XIIth International Conference on AIDS and STDs in Africa, Ouagadougou, Burkina Faso, 12/9-13, Session 12DT3-2.

Dolmans, W. M. V., A. M. Van Loon, R. Van den Akker, et al. 1989 Prevalence of HIV-1 Antibody among Groups of Patients and Healthy Subjects from a Rural and Urban Population in the Mwanza ... AIDS, vol. 3, no. 5, pp. 297-299.

Daley, C. L., L. L. Chen, P. M. Small, et al. 1992 Pulmonary Complications of HIV Infection in Tanzania American Review of Respiratory Disease, vol. 145, p. A821.

Hemed, Y., J. Minja, E. Naegele, et al. 1989 Seroepidemiological Study of Sexually Transmitted Agents (HIV, HBV, T. Pallidum) among Patients in the Mbeya Region (South Ta. ... IV Internat. Conf.: AIDS and Assoc. Cancers in Africa, Marseille, Oct. 18-20, Abstract 080.

Haukenes, G., J. Shao, F. Mhalu, et al. 1992 The AIDS Epidemic in Tanzania: Rate of Spread of HIV in Blood Donors and Pregnant Women in Dar es Salaam Scandinavian Journal of Infectious Diseases, vol. 24, pp. 701-706.

Haji, S. H., A. K. Ali, A. A. Othman, et al. 1995 Prevalence of HIV Infection in Inmate Prisoners Pemba Island, Zanzibar, Tanzania IX International Conference on AIDS and STD in Africa, Kampala, Uganda, 12/10-14, Abstract TuD128.

Hoffmann, O., L. Maboko, B. Wolff, et al. 2002 Early Infection with HIV-1 among Women in a High-Risk Environment in Mbeya, Tanzania XIV International AIDS Conference, Barcelona, Spain, 7/7-12, Abstract MoPeC3509.

Johns Hopkins University 1986 AIDS -- A Public Health Crisis Population Information Program, Population Reports, Issues in World Health, July-Aug., Series L, no. 6, pp. 194-228.

Jacobson, M. L.; N. King'Ori, E. Ngoga, et al. 1993 HIV1 Prevalence in a Nomadic Population in Tanzania VIII International Conference on AIDS in Africa, Marrakech, Morocco, 12/12-16, Poster T.P.C.095.

Kipuyo, E. L. 1997 HIV-Trends and Behaviour Changes in Tanzania Presented at the UNAIDS Regional Workshop on "Evidence of Behavioural Change in the Context of HIV Decline in Uganda", 10 - 13 February, Nairobi, Kenya.

Kwesigabo, G., J. Z. J. Killewo, W. Urassa, et al. 2000 Monitoring of HIV-1 Infection Prevalence and Trends in the General Population Using Pregnant Women as a Sentinel ... Journal of Acquired Immune Deficiency Syndromes, vol. 23, no. 5, pp. 410-417.

Kilewo, C., A. Massawe, E. Lyamuya, et al. 2001 HIV Counseling and Testing of Pregnant Women in Sub-Saharan Africa: Experiences from a Study on Prevention ... Journal of Acquired Immune Deficiency Syndromes, vol. 28, no. 5, pp. 458-462.

Kapiga, S. H., D. Mwakagile, D. Spiegelman, et al. 2000 Predictors of CD4+ Lymphocyte Count among HIV-Seropositive and HIV-Seronegative Pregnant Women in Dar es Salaam, Tanzania East African Medical Journal, vol. 77, no. 4, pp. 206-211.

Lisekie, F., H. Grosskurth, A. Klokke, et al. 1992 Comparison of Sentinel and Cross-Sectional Study Data: Is Sentinel Surveillance a Useful Tool to Monitor the HIV/AIDS Epidemic? VII International Conference on AIDS in Africa, Yaounde, Cameroon, 12/8-11, Poster T.P.035.

Lyamuya, E. F., E. Urassa, E. Mbena, et al. 1996 Stabilization of HIV-1 Seroprevalence in Pregnant Women in Dar es Salaam, Tanzania: Towards Reaching a Plateau Phase? XI International Conference on AIDS, Vancouver, 7/7-14, Poster Tu.C.2463.

Mhalu, F., U. Bredberg-Raden, E. Mbena, et al. 1987 Prevalence of HIV Infection in Healthy Subjects and Groups of Patients in Tanzania AIDS, vol. 1, no. 4, pp. 217-221.

Mhalu, F., M. Ngaiza, A. B. M. Swai, et al. 1991 Factors Associated with HIV Infection among a Sample of Bar and Restaurant Workers in Dar es Salaam, Tanzania VII International Conference on AIDS, Florence, Italy, 6/16-21, Poster W.C.3087.

Mwakagile, D. S. M., A. B. M. Swai, S. Mwambimanga, et al. 1991 Socio-Epidemiological and Microbiological Aspects of Sexually Transmitted Diseases and HIV Infection in Dar es Salaam, Tanzania VI International Conference on AIDS in Africa, Dakar, Senegal, 12/16-19, Abstract W.A.144.

Matasha, E., J. Chungalucha, H. Grosskurth, et al. 1992 Commercial Sexual Workers Intervention Programme: A Pilot Project in Northern Tanzania: Operational Data and International. ... VIII International Conference on AIDS, Amsterdam, 7/19-24, Poster PoD 5639.

Mwakagile, D. S. M., A. B. M. Swai, K. J. Pallangyo, et al. 1992 Trend of Anogenital Warts among Patients Seen at a Referral Clinic for Sexually Transmitted Diseases in Dar es Salaam, Tanzania VII International Conference on AIDS in Africa, Yaounde, Cameroon, 12/8-11, Poster W.P.190.

Mhalu, F., A. Swai, D. Mwakagile, et al. 1992 Surveillance and Control of HIV-1 Transmission among Female Bar workers in Dar es Salaam 1986-1991 VII International Conference on AIDS in Africa, Yaounde, Cameroon, 12/8-11, Poster T.P.108.

Mwakagile, D., A. Swai, J. Kitange, et al. 1993 Epidemiology of STDs in Dar es Salaam, Tanzania VIII International Conference on AIDS in Africa, Marrakech, Morocco, 12/12-16, Session M.O.P.058.

Mwakagile, D., R. Mkuna, A. B. M. Swai, et al. 1993 Diagnosis of Sexually Transmitted Diseases (STDs) in Pregnant Women VIII International Conference on AIDS in Africa, Marrakech, Morocco, 12/12-16, Abstract M.O.P.027.

Mwambingu, S. H., G. Rutahindurwa, F. Mhalu, et al. 1995 Development of STD Services in an Urban Clinic in Dar es Salaam, Tanzania IX International Conference on AIDS and STD in Africa, Kampala, Uganda, 12/10-14, Poster MoB452.

Msamanga, G. I., E. Urassa, D. Spiegelman, et al. 1996 Socioeconomic Status and Prevalence of HIV Infection among Pregnant Women in Dar es Salaam, Tanzania XI International Conference on AIDS, Vancouver, 7/7-14, Poster Tu.C.2464.

Mlika-Cabanne, N., M. Brauner, F. Mugusi, et al. 1995 Radiographic Abnormalities in Tuberculosis and Risk of Coexisting HIV Infection: Results from Dar-es-Salaam, Tanzania, and . . American Journal of Respiratory and Critical Care Medicine, vol. 152, no. 2, pp. 786-793.

Mwakagile, D., A. B. M. Swai, E. Sandstrom, et al. 1996 High Frequency of Sexually Transmitted Diseases among Pregnant Women in Dar es Salaam, Tanzania: Need for Intervention East African Medical Journal, vol 73, no. 10, pp. 675-679.

Mwakagile, D., E. Mmari, C. Makwaya, et al. 1998 Unusual Characteristics of Adolescents and Youths Attending a New Special STD Clinic for Them in Dar es Salaam; . . 12th World AIDS Conference, Geneva, 6/28 - 7/3, Abstract 22203.

Mayaud, P., D. K. Gill, H. A. Weiss, et al. 2001 The Interrelation of HIV, Cervical Human Papillomavirus, and Neoplasia among Antenatal Clinic Attenders in Tanzania Sexually Transmitted Infections, vol. 77, no. 4, pp. 248-254.

Ministry of Health 2000 HIV/AIDS/STDs Surveillance Report Ministry of Health and Social Welfare, Zanzibar AIDS Control Programme, no. 2, report.

Nkya, W. M. M. M., W. P. Howlett, C. Assenga, et al. 1987 Seroepidemiology of HIV-1 Infection in the Kilimanjaro Region AIDS and Associated Cancers in Africa, Naples, pp. 61-65.

Nkya, W. M. M. M., W. P. Howlett, C. Assenga, et al. 1988 AIDS Situation in Northern Tanzania III International Conference: AIDS and Associated Cancers in Africa, Sept. 14-16, Poster TP 9.

Nkya, L., B. F. Lyamuya, A. Outwater, et al. 1993 The Pattern of Sexually Transmitted Diseases in a Group of Commercial Sex Workers in Morogoro, Tanzania VIII International Conference on AIDS in Africa, Marrakech, Morocco, 12/12-16, Poster T.P.C.083.

Noronha, D., K. J. Pallangyo, B. N. Ndosu, et al. 1991 Radiological Features of Pulmonary Tuberculosis in Patients Infected with Human Immunodeficiency Virus East African Medical Journal, vol. 68, no. 3, pp. 210-215.

Othman, A. A., A. M. Shein, M. I. Hassan, et al. 1993 HIV Seroprevalence in Patients Attending STD Clinic at Mnazi Mmoja Hospital (MMH), Zanzibar VIII International Conference on AIDS in Africa, Marrakech, Morocco, 12/12-16, Poster Th.P.C.075.

Ocheng, D., A. Msauka, C. Mbuya, et al. 1993 Prevalence of Sexually Transmitted Diseases (STDs) in Females at 7 Truck Stops in Tanzania VIII International Conference on AIDS in Africa, Marrakech, Morocco, 12/12-16, Poster Th.P.C.070.

Othman, A. A., A. M. Shein, A. K. Ali 1994 Laboratory Findings for the Prevalence of HIV and Other STDs in Antenatal Attendees at Mnazi Mmoja Hospital: A Pilot Study Tenth International Conference on AIDS, Yokohama, Japan, 8/7-12, Abstract P.A.0190.

Petry, K. U., H. Kingu 1996 HIV Infection among Pregnant Women in Lindi, Tanzania, 1989-1993 International Journal of STD and AIDS, vol. 7, pp. 265-268.

Riedner, G., Y. Hemed, F. Minja, et al. 1993 The Use of Serologic Trends of HIV and Syphilis for the Evaluation of the Mbeya Regional ACP Tanzania 1986-1992 IX International Conference on AIDS, Berlin, 6/6-11, Poster PO-C29-3263.

Richter, C., M. J. W. Koelemay, A. B. M. Swai, et al. 1995 Predictive Markers of Survival in HIV-Seropositive and HIV-Seronegative Tanzanian Patients with Extrapulmonary Tuberculosis Tubercle and Lung Disease, vol. 76, no. 6, pp. 510-517.

Seif, M. J., S. H. Haji, A. M. Shein, et al. 1993 Anonymous Unlinked Testing HIV Prevalence in Pregnant Mothers in Pemba (Zanzibar) VIII International Conference on AIDS in Africa, Marrakech, Morocco, 12/12-16, Abstract Th.P.C.076.

Shushu, M. L., J. Chagalucha, J. Ngalula, et al. 2001 Prevalence of STIs in Women Attending Gynaecology Clinics in Mwanza, Tanzania Presented at International Congress of Sexually Transmitted Infections, ISSTD/ISTDI, Berlin, Germany, 24-27 June 2001, Abstract in International Journal of STD & AIDS, vol. 12, suppl. 2, p. 91.

Ter Meulen, J., K. M. Wittkowski, J. J. Kidenya, et al. 1989 Evaluation of Seroepidemiological Associations between HIV-Infection, Hepatitis B and Other Sexually Transmitted Diseases in. . European Journal of Epidemiology, vol. 5, no. 2, pp. 158-163.

Tanzania Ministry of Health 1992 National AIDS Control Programme Surveillance Report No. 7., December, Epidemiology Unit, NACP.

Tanzania Ministry of Health 1994 National AIDS Control Programme Surveillance Report, no. 8, June, Epidemiology Unit, NACP.

Tanzania Ministry of Health 1995 National AIDS Control Programme HIV/AIDS/STD Surveillance Surveillance report no. 9, December 1994, Epidemiology Unit, NACP.

Tanzania Ministry of Health 1996 National AIDS Control Programme - HIV/AIDS/STD Surveillance Surveillance report no. 10, December 1995, Epidemiology Unit, NACP.

Tanzania Ministry of Health 1997 National AIDS Control Programme - HIV/AIDS/STDs Surveillance Surveillance report, no. 11, December 1996, Epidemiology unit, NACP.

Tanzania Ministry of Health 1998 National AIDS Control Programme - HIV/AIDS/STDs Surveillance Surveillance report, no. 12, December 1997, Epidemiology unit, NACP.

Tanzania Ministry of Health 1999 National AIDS Control Programme - HIV/AIDS/STDs Surveillance Surveillance Draft report, no. 13, December 1998, Epidemiology unit, NACP.

Tanzania Ministry of Health 2000 National AIDS Control Programme - HIV/AIDS/STD Surveillance Surveillance report, no. 14, December 1999, Epidemiology unit, NACP.

Tanzania Ministry of Health 2001 National AIDS Control Programme - HIV/AIDS/STD Surveillance Surveillance report, no. 15, January - December 2000, NACP.

Tanzania Ministry of Health 2002 Surveillance of HIV and Syphilis among Antenatal Clinic Enrollees: 2001-2002 National AIDS Control Programme, Ministry of Health, Dar es Salaam, Tanzania, unpublished report.

Urassa, E., F. S. Mhalu, E. Mbena, et al. 1990 Prevalence of HIV-1 Infection among Pregnant Women in Dar es Salaam, Tanzania V International Conference: AIDS in Africa, Kinshasa, Zaire, Oct. 10-12, Poster T.P.E.22.

Van Cleeff, M. R. A., H. J. Chum 1992 The Influence of HIV on the Epidemiology of Tuberculosis: An Analysis Made on Basis of a Pilot Study in Tanzania VIII International Conference on AIDS, Amsterdam, 7/19-24, Poster PoC 4033.

Van Den Broek, J., M. W. Borgdorff, N. G. Pakker, et al. 1993 HIV-1 Infection as a Risk Factor for the Development of Tuberculosis: A Case-Control Study in Tanzania International Journal of Epidemiology, vol. 22. no. 6, pp. 1159-1165.

Watson-Jones, D., J. Chagalucha, B. Gumodoka, et al. 2002 Syphilis in Pregnancy in Tanzania. I. Impact of Maternal Syphilis on Outcome of Pregnancy Journal of Infectious Diseases, vol. 186, no. 7, pp. 940-947.

Watson-Jones, D., B. Gumodoka, H. Weiss, et al. 2002 Syphilis in Pregnancy in Tanzania. II. The Effectiveness of Antenatal Syphilis Screening and Single-Dose Benzathine ... Journal of Infectious Diseases, vol. 186, no. 7, pp. 948-957.

**Websites:**     [www.aids.africa.com](http://www.aids.africa.com)

## Annex: HIV surveillance by site

Group	Area	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003		
Pregnant women	Outside major urban areas										2.80			0.60	0.93					
	Chake chake, Pemba																			
	Chimalala, Mbeya region/ rural		4.17	6.25	8.80	9.50	8.00	10.79	16.00	10.50	17.00	15.90	12.50	12.10	15.20			17.05		
	Dodoma region		3.00																6.08	
	Handali, Dodoma region/ rural																		0.49	
	Hedaru, Kilimanjaro region/ rural																			5.44
	Igekemaja, Mwanza region															4.20				
	Igoma, Mwanza region															13.80				
	Ilembo, Mbeya region/ rural																			7.11
	Ipogoro, Iringa region/ urban															32.10				
	Iringa Regional Hospital, Iringa re													24.90	40.10	4.60				
	Isoko, Mbeya region/ rural		2.94	2.00	2.40	6.60	18.00	8.47	8.00	10.30	7.50	8.10	10.20	19.10	13.50					
	Ilete, Mbeya region/ rural		1.72	9.09	6.40	3.90	5.26	15.52	5.10	14.80	5.50	14.80	11.80	11.60	23.30					
	Katoro, Kagera district/ semiurban																			3.00
	Kibaigwa, Dodoma region/ rural																			10.50
	Kimeya, Kagera district/ rural																			6.08
	Kiseso, Mwanza region															10.80				
	Kivunge, Unguja										6.30	0	1.90		0.50	0.96				
	Kiwanjampaka, Mbeya region/ urban		7.00	10.64				17.00	22.34	19.50		17.00	22.50	20.50	23.00	23.30				17.96
	Korogwe, Tanga region															9.30				
	Kyela, Mbeya region/ rural			21.21	14.60	17.50	30.36	27.23	27.50	32.90	25.90	25.00	24.00	29.50	21.60					17.16
	Ligula, Mtwana region/ urban																			12.46
	Lindi district			0.44					8.68											
	Madaba, Ruvuma region							6.60	12.90											
	Mafinga, Iringa region						21.03	25.04							20.90	16.80				
	Majengo, Kilimanjaro region/ urban																			7.55
	Makole, Dodoma region/ urban																			9.70
	Makongoro, Mwanza region															16.40				
	Makorora, Tanga region															8.55				
	Makunduchi, Unguja												0							
	Mangaka, Mtwara region/ semiurban																			4.30
	Masama, Kilimanjaro region/ rural																			2.26
	Mbozi, Mbeya region									15.00	13.90	17.00	24.00			19.00				
Meta, Mbeya region/ urban		10.28	16.92				25.00	13.71	16.00		14.60	17.90	12.50	13.50	17.00					
Micheweni, Pemba										8.90	0			1.10	0					
Mkoani, Pemba											5.40									
Mkula (Magu), Mwanza region/ urban						3.70	4.60	5.40		5.40										
Mlalo, Tanga region															2.70					
Mnazi Mmoja Hospital, Zanzibar area							0.50	2.29												
Mnazimmoja, Unguja										2.40	0.20	1.60		0.80	0.75					
Morogoro Regional Hospital, Morogor														18.40						
Musoma, Mara region						5.90	6.50	7.70												
Mwambani, Mbeya region/ rural			12.00	8.50	12.90	8.00	10.73	13.00	17.50	16.00	13.70	14.50	11.00	13.00						
Mwanjelwa, Mbeya region/ urban		11.00	7.29				11.00	23.25	19.60			36.00								

Group	Area	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003		
Pregnant women	Outside major urban areas	Mwanza region, Mwanza							15.27							15.00				
		Namanyere, Rukwa region/ rural						11.33	8.33	19.00	11.20		11.20							
		Namtumbo, Ruvuma region/ rural						3.53	6.74	3.20	5.60		4.00							
		Nanguruwe, Mtwara region					4.44				0									
		Nanyamba, Mtwara region/ rural																	4.00	
		Nkwenda, Kagera district/ rural																		3.88
		Nyasio, Mara region					9.00	7.01	8.00		7.20									
		Pemba island				0.29			0.60	0.77	0.30									
		Ruanda, Mbeya region/ urban											24.00	18.10	18.80	17.50	20.70			
		Rural sites, Mwanza region/ rural					3.70	3.90	5.49	5.46	5.38									
		Shinyanga region									10.90									
		Singida region		3.60																
		Songea, Ruvuma region/ urban							9.67	16.13	15.70	14.20		11.00						
		Sumbawanga, Rukwa region/ urban							12.00	23.18	13.30	22.20		21.00						
		Tandahimba, Mtwara region/ semiurba																		3.92
		Tanga region		7.00																
		Turiani DDH, Morogoro region/ rural													9.80					
		Umbwe, Kilimanjaro region/ rural						2.34	6.37	15.84		0	9.10	10.00	19.93	19.20	16.60			8.77
		Unguja & Pemba Islands, Zanzibar ar		0.26	0.53	0.63	0.61	0.68									0.68			
		Urban sites, Mbeya region/ urban					12.20	15.30				20.70								
		Urban sites, Mwanza region/ urban			7.96	11.66	12.31	11.19	10.03	12.33	11.70	8.78								
		Welama-songa, Mwanza region															5.00			
		Wete, Pemba											0							
		3 hospitals, Mwanza region		6.02											10.27	4.00				
		5 rural dispensaries, Arusha region								0.98										
		Arusha region			5.50															
		Bahi, Dodoma region/ rural																		1.16
		Bukoba					22.24	20.04	27.72	17.29	17.30		13.00			7.00	12.50			8.33
		4 district hospitals, Dar es Salaam										13.70								
		Buguruni, Dar es Salaam region																		16.29
		Dar-es-Salaam (1), Dar es Salaam			7.80					16.10			12.99	13.73						
		Kasorobo, Dar es Salaam region																		10.00
		Kasorobo-Temeke, Dar es Salaam regi														15.30	14.30			
Kigamboni, Dar es Salaam region																		11.98		
Kigamboni-Temeke, Dar es Salaam reg														14.10	10.10					
Magomeni HC ANC, Dar es Salaam								15.18												
Muhimbili Medical Center, Dar es Sa				8.93	8.95	9.12	11.03	10.61	13.81	12.17										
Oysterbay, Dar es Salaam region																		10.94		
Sinza-Kinondoni, Dar es Salaam regi														18.10						
Temeke, Dar es Salaam								15.30		7.30										
Temeke, M'Nya mala, A.K. Clinic, Dar						10.33														
Sex workers	Major urban areas		35.71	43.73	44.97	42.86		49.46												

Group	Area		1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003
Sex workers	Major urban areas	Dar-es-Salaam (2), Dar es Salaam		42.08	26.71														
		Outside major urban areas	Arusha region		18.60														
		Bulyanhulu														44.60			
		Dodoma region		19.50															
		Geita																39.00	
		Kagera region																	
		Kilimanjaro region, Kilimanjaro reg	14.80																
		Mbeya region																	68.00
		Morogoro								60.87									
		Moshi region		31.90															
	Mwanza town							39.81											
	Singida region		7.50																
	Tanga region		25.70																
Injecting drug users																			
STI patients	Major urban areas	Dar-es-Salaam (1), Dar es Salaam	20.72	24.81	33.99	22.77	18.70						14.75			46.00	41.50		
		Dar-es-Salaam (2), Dar es Salaam					29.44												
		Mnazi Mmoja & Magomeni, Dar es Sala					16.80												
		Muhimbili Medical Center, Dar es Sa							23.97										
	Outside major urban areas	Chimala, Mbeya region/ rural												18.00					
		Isoko, Mbeya region/ rural												22.00					
		Itete, Mbeya region/ rural												27.00					
		Kiwanjampaka, Mbeya region/ urban												38.00					
		Kyela, Mbeya region/ rural												25.00					
		Mbeya region		22.50	22.48	27.50	32.00	34.30											
		Mbozi, Mbeya region												43.00					
		Mnazi Mmoja Hospital, Zanzibar area							5.16										
		Mwambani, Mbeya region/ rural													36.00				
		Referral Hospital, Mbeya region/ ur													17.00				
		Ruanda Municipal, Mbeya region/ urb													36.00				
		St. Joseph Hospital, Kagondo (rural)	33.34																
Men having sex with men		Major urban areas	Ilala district, Dar es Salaam regio											52.37					
	Kinondoni district, Dar es Salaam r													53.14					
	Muhimbili Medical Center, Dar es Sa			35.83			65.62	55.22			54.79								
	Temeke district, Dar es Salaam regi													52.37					
	Outside major urban areas	Arusha region					23.51							24.53					
		Coast region												51.60					
		Dodoma region							21.67					38.63					
		Iringa region								59.04									
		Kagera region					36.47							37.96					
		Kigoma region								19.55									
		Kilimanjaro region					42.68							45.37					
		Lindi region							13.33						31.52				
		Mara region								17.90					29.24				
		Mbeya region						53.39							70.54				
		Mbeya region, Mbeya region/ urban				54.00													
		Mnazimmoja, Unguja															32.00		
Morogoro region							33.00						32.98						
Mwanza region, Mtwara region								9.62					27.07						
Mwanza region, Mwanza region/ rural						24.62							34.14						

Group	Area	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	
Tuberculosis patients	Outside major urban areas	Mwanza region, Mwanza region/urban				54.05													
		National Reference Hospital for TB,	6.34																
		Pwani region						30.40				51.58							
		Rukwa region							44.97										
		Ruvuma region							39.42										
		Shinyanga region							26.30				31.15						
		Singida region								20.67			32.17						
		Tabora region								16.48			47.83						
		Tanga region					19.73						43.31						
		Temeke district								46.20									
		Zanzibar region											21.55						